

# Air Pollution: Health Risk Assessment Mount Maunganui

---

1 June 2023

**PREPARED FOR:** Toi Te Ora Public Health  
**CLIENT REPORT No:** FW23019

# ACKNOWLEDGEMENTS

---

The author wishes to acknowledge Bay of Plenty Regional Council who assisted with provision of ambient air quality monitoring data for this assessment. Internal, external and international peer and expert reviewers who assessed this report are also gratefully acknowledged.

## DISCLAIMER

This report has been released to Toi Te Ora.

The Institute of Environmental Science and Research Limited (ESR) has used all reasonable endeavours to ensure that the information contained in this client report is accurate. However, ESR does not give any express or implied warranty as to the completeness of the information contained in this client report or that it will be suitable for any purposes other than those specifically agreed by ESR and Toi Te Ora in writing.



# CONTENTS

---

<b>EXECUTIVE SUMMARY</b> .....	<b>1</b>
<b>1. INTRODUCTION</b> .....	<b>1</b>
1.1 BACKGROUND .....	1
1.2 PRINCIPLES OF ASSESSMENT .....	6
<b>2. HEALTH EFFECTS OF AIR CONTAMINANTS</b> .....	<b>8</b>
2.1 PARTICULATE MATTER (PM <sub>10</sub> AND PM <sub>2.5</sub> ).....	8
2.2 NITROGEN DIOXIDE (NO <sub>2</sub> ) .....	9
2.3 SULPHUR DIOXIDE (SO <sub>2</sub> ) .....	10
2.4 HYDROGEN SULPHIDE (H <sub>2</sub> S).....	11
2.5 BENZENE .....	13
2.6 ODOUR .....	15
<b>3. METHODOLOGY</b> .....	<b>16</b>
3.1 CONTAMINANTS OF PUBLIC HEALTH CONCERN.....	17
3.2 QUANTIFIED RISK ASSESSMENT: PM <sub>10</sub> , PM <sub>2.5</sub> AND NO <sub>2</sub> .....	18
3.3 QUALIFIED RISK ASSESSMENT: SO <sub>2</sub> , H <sub>2</sub> S, BENZENE AND ODOUR .....	24
<b>4. QUANTIFIED CHRONIC EXPOSURE: PM<sub>10</sub>, PM<sub>2.5</sub> &amp; NO<sub>2</sub> IN MOUNT MAUNGANGUI</b> .....	<b>25</b>
4.1 PM <sub>10</sub> .....	25
4.2 PM <sub>2.5</sub> .....	27
4.3 NO <sub>2</sub> .....	28
<b>5. QUANTIFIED HEALTH IMPACTS: PM<sub>10</sub>, PM<sub>2.5</sub> &amp; NO<sub>2</sub> IN MOUNT MAUNGANGUI..</b>	<b>30</b>
5.1 PM <sub>2.5</sub> AND NO <sub>2</sub> .....	30
5.2 PM <sub>10</sub> .....	31
5.3 MEETING THE WHO AQG 2021 GUIDELINES .....	31
5.4 SENSITIVITY TESTING .....	32
5.5 UNCERTAINTY & CONFIDENCE ASSESSMENT .....	33
5.6 MODEL LIMITATIONS .....	35
<b>6. QUALITATIVE ASSESSMENT: SO<sub>2</sub>, H<sub>2</sub>S, BENZENE &amp; ODOUR IN MOUNT MAUNGANGUI.....</b>	<b>36</b>
6.1 SULPHUR DIOXIDE.....	36
6.2 HYDROGEN SULPHIDE .....	38
6.3 BENZENE .....	40
6.4 ODOUR .....	43
<b>7. CONCLUSIONS</b> .....	<b>44</b>

# CONTENTS CTD

---

REFERENCES .....	47
APPENDIX A: MT MAUNGANUI SO <sub>2</sub> DATA.....	51

## LIST OF TABLES

TABLE 1:	NEW ZEALAND AND WHO AIR QUALITY GUIDELINES AND STANDARDS FOR PARTICULATE MATTER.....	9
TABLE 2:	NEW ZEALAND AND WHO QUALITY GUIDELINES AND STANDARDS FOR NO <sub>2</sub> .....	10
TABLE 3:	NEW ZEALAND AND WHO SHORT-TERM AIR QUALITY GUIDELINES AND STANDARDS FOR SO <sub>2</sub> .....	11
TABLE 4:	ACUTE HEALTH EFFECTS OF H <sub>2</sub> S AT VARYING CONCENTRATIONS [SOURCE: ESR 2022].....	13
TABLE 5:	COMPARISON OF MEAN 2019 PM <sub>10</sub> CONCENTRATIONS AT MOUNT MAUNGANUI MONITORING LOCATIONS.....	26
TABLE 6:	MEAN 2019 PM <sub>10</sub> CONCENTRATIONS ASSIGNED TO EACH CENSUS AREA UNIT .....	27
TABLE 7:	MEAN 2019 PM <sub>2.5</sub> CONCENTRATIONS ASSIGNED TO EACH CENSUS AREA UNIT .....	28
TABLE 8:	MEAN 2019 NO <sub>2</sub> CONCENTRATIONS ASSIGNED TO EACH CENSUS AREA UNIT.....	29
TABLE 9:	ESTIMATED PREMATURE MORTALITY AND MORBIDITY ASSOCIATED WITH ANNUAL PM <sub>2.5</sub> AND NO <sub>2</sub> .....	30
TABLE 10:	ESTIMATED PREMATURE MORTALITY ASSOCIATED WITH ANNUAL PM <sub>10</sub> .....	31
TABLE 11:	ESTIMATED AVOIDED HEALTH IMPACTS AND SOCIAL COSTS IF WHO GUIDELINES MET IN OTŪMOETAI AND MOUNT MAUNGANUI VERSUS BASE CASE.....	32
TABLE 12:	SENSITIVITY OF THE HAPINZ 3 SOCIAL COST ESTIMATES FOR PM <sub>2.5</sub> AND NO <sub>2</sub> TO DIFFERENT PARAMETERS VERSUS THE BASE CASE (IN 2019 NZD).....	32
TABLE 13:	SUMMARY HOURLY H <sub>2</sub> S CONCENTRATIONS AT WHAREROA MARAE 2016 – 2020 .....	40
TABLE 14:	HOURLY BENZENE CONCENTRATIONS MEASURED AT MOUNT MAUNGANUI, 2020-2021 .....	42

# CONTENTS CTD

---

## LIST OF FIGURES

FIGURE 1: MT MAUNGANUI AIRSHED (RED OUTLINE) AND ADJACENT RESIDENTIAL AREAS (ORANGE OUTLINE APPROX. 2013 CAUS) .....	3
FIGURE 2: ACTIVITIES SENSITIVE TO AIR POLLUTION IN AND AROUND THE MOUNT MAUNGANUI AIRSHED .....	4
FIGURE 3: BAY OF PLENTY REGIONAL COUNCIL AMBIENT AIR QUALITY MONITORING FOR PM <sub>10</sub> AND SO <sub>2</sub> IN THE MT MAUNGANUI AIRSHED.....	5
FIGURE 4: OVERVIEW OF AN AIR POLLUTION HEALTH RISK ASSESSMENT PROCESS.....	16
FIGURE 5: ANNUAL TOTAL SUSPENDED PARTICULATE AND PM <sub>10</sub> MEASURED IN OTŪMOETAI AND SELECT MT MAUNGANUI INDUSTRIAL LOCATIONS ..	20
FIGURE 6: NZ STATS (2020) WHERE WE LIVE VERSUS WHERE WE WORK (USING 2018 CENSUS DATA) .....	23
FIGURE 7: NATIONAL MORTALITY RATES 2009-2018.....	24
FIGURE 8: 2019 PM <sub>10</sub> MONITORING LOCATIONS AND ANNUAL PM <sub>10</sub> CONCENTRATIONS MEASURED IN THE MT MAUNGANUI AIRSHED .....	26
FIGURE 9: HOURLY H <sub>2</sub> S CONCENTRATIONS MEASURED AT WHAREROA MARAE; 26 SEP 2015 – 14 OCT 2020 .....	39
FIGURE 10: BENZENE SOURCES MOUNT MAUNGANUI (TRAFFIC, FUEL TANKS, OIL RE-REFINERY) IN RELATION TO WHAREROA MARAE & TAURANGA BRIDGE MARINA .....	41

# EXECUTIVE SUMMARY

---

This report is an air pollution health risk assessment for the Mount Maunganui area, prepared in response to a request by Toi Te Ora Public Health. The intent is to provide information to polluters, regulatory agencies, and the affected community on the potential scale of adverse health outcomes from existing air quality with the aim of working together to reduce discharges.

The approach taken has been to qualitatively and, where practicable, quantitatively describe and assess potential health risks of exposure to identified air pollutants in the Mount Maunganui area in accordance with good practice (WHO, 2014). The qualitative assessment reviews available data for the period ending 31 December 2021. The quantitative assessment uses a base year of 2019, which pre-dates potential impacts of the COVID-19 pandemic on ambient air quality.

The pollutants assessed were:

- Particulate matter (PM) less than 10 micrometres ( $\mu\text{m}$ ) in diameter ( $\text{PM}_{10}$ ) and less than 2.5  $\mu\text{m}$  in diameter ( $\text{PM}_{2.5}$ )
- Nitrogen dioxide ( $\text{NO}_2$ )
- Sulphur dioxide ( $\text{SO}_2$ )
- Hydrogen sulphide ( $\text{H}_2\text{S}$ )
- Benzene ( $\text{C}_6\text{H}_6$ )
- Odour

Toi Te Ora Public Health also requested assessment of polycyclic aromatic hydrocarbons (PAHs) but this was unable to be undertaken due to a lack of ambient air quality monitoring data. Methyl bromide was excluded from consideration.

First, we briefly described the potential health effects of these pollutants with reference to the most up to date state of knowledge on the science of the health effects of key pollutants ( $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$  and  $\text{NO}_2$ ) in New Zealand (Hales et al., 2021) and globally (WHO, 2022). Next, we reviewed available ambient air quality monitoring data in Mount Maunganui and surrounding areas. This showed that concentrations of some air contaminants are elevated in some locations relative to health and wellbeing-based ambient criteria.

Our assessment was in two parts – quantitative and qualitative. It should be noted that our assessment addresses only identified pollutants, individually, which excludes consideration of potential synergistic or cumulative impacts. It also does not consider wider impacts including positive social benefits arising from the sources of discharges to air (for example, contribution to the local and national economy).

## Quantitative Assessment

The quantitative assessment focused on long-term exposure which, for PM and  $\text{NO}_2$ , is known to result in an order of magnitude greater significance to public health than acute exposure (WHO, 2021). Our assessment was comparative, using current monitoring of annual average concentrations as a chronic exposure metric.

Specifically, we compared public exposure to key pollutants (PM<sub>10</sub>, PM<sub>2.5</sub> and NO<sub>2</sub>) in Mount Maunganui which is immediately adjacent to the Port and industrial area, with public exposure in Otūmoetai, a residential suburb of Tauranga around five kilometres away. The health effects of these pollutants (only) have recently been quantified for New Zealand (Hales *et al.*, 2021, Kuschel *et al.*, 2022).

Our approach to estimating chronic population exposure to key pollutants was one of caution to avoid overstating potential impacts. This means the estimates are not conservative and likely underestimate effects.

The modelling estimates show that, compared with Otūmoetai, in Mount Maunganui there were:

- Around five additional premature deaths in adults (>30 years) each year associated with exposure to long-term concentrations of **PM<sub>2.5</sub> and NO<sub>2</sub>**. For context, the total mortality from all non-external causes<sup>1</sup> in Mount Maunganui for the year 2019 was 145 so this estimate represents around 3% of deaths in that year.
- An additional four cardiovascular and six respiratory hospitalisations per year associated with long-term exposure to PM<sub>2.5</sub> and NO<sub>2</sub>.
- An additional 1,256 restricted activity days per year across the population associated with long-term exposure to PM<sub>2.5</sub>.
- Two additional cases of asthma per year in under 18-year-olds associated with long-term exposure to NO<sub>2</sub>.
- Estimated social costs due to additional mortality and morbidity of \$22 million (NZ\$2019).

A statistical sensitivity analysis estimates that:

- The number of additional premature deaths for adults (>30 years) per year associated with long-term exposure to PM<sub>2.5</sub> and NO<sub>2</sub> in Mount Maunganui as compared with Otūmoetai has a 95% confidence interval of 3 to 6 (relative to the base case of 5 deaths).
- The low and high social cost estimates associated with long-term exposure to PM<sub>2.5</sub> and NO<sub>2</sub> in Mount Maunganui as compared with Otūmoetai are between \$19 million and \$29 million per year (relative to the base case of \$22 million, all in NZ\$2019).

Single-pollutant modelling for **PM<sub>10</sub>** (only) estimated that the Mount Maunganui area had 13 additional premature deaths in adults (> 30 years) each year (95% confidence interval 11 to 15) when compared with Otūmoetai. This estimate represents around 9% of total mortality from all non-external causes in Mount Maunganui that year, which is higher than the estimates associated with long-term exposure to PM<sub>2.5</sub> and NO<sub>2</sub>. It should be noted that the PM<sub>10</sub> modelling is not additive to the estimate of effects associated with PM<sub>2.5</sub> and NO<sub>2</sub>, rather it is a separate estimate.

At Toi Te Ora Public Health's request, we modelled the hypothetical scenario of all areas, including Otūmoetai, meeting the annual WHO 2021 global air quality guidelines for PM<sub>2.5</sub> (5 µg/m<sup>3</sup>) and NO<sub>2</sub> (10 µg/m<sup>3</sup>). This would have a modest impact in averting two premature deaths per year with an associated averted social cost of \$11 million (NZ\$2019) due to hypothetical reductions in annual PM<sub>2.5</sub> (as our assessment assumed all areas already meet the annual WHO guideline for NO<sub>2</sub>).

---

<sup>1</sup> i.e., deaths excluding accidents and violence

Our assessment of uncertainty concludes there is a moderate degree of confidence in the modelling estimates.

## Qualitative Assessment

### *Sulphur Dioxide*

Following complaints of adverse health effects by residents', Bay of Plenty Regional Council (BOPRC) commenced ambient air quality monitoring at Whareroa Marae in 2015. This revealed breaches of the national environmental standards for SO<sub>2</sub> caused by a neighbouring fertiliser works. Regulatory action by BOPRC, coupled with mitigation by the fertiliser works, significantly reduced emissions and there have been no breaches of national environmental standards for SO<sub>2</sub> since 2016. The elevated acute (10-minute) SO<sub>2</sub> concentrations may have resulted in some short-term, transient effects (bronchoconstriction, nose or throat irritation) at nearby locations such as Whareroa Marae.

In late 2018 BOPRC commenced monitoring for SO<sub>2</sub> at five other locations of the Mount Maunganui Airshed. In 2019 there were occasional 10-minute and hourly concentrations that could cause asthma and adverse respiratory effects with multiple exceedances of WHO guidelines and national environmental standards for SO<sub>2</sub> at these monitoring stations. However, from 1 January 2020, the short-term peaks disappeared at most monitoring locations. This date coincides with reductions in ship emissions due to the implementation of MARPOL Annex VI from this date.

The exceptions were Whareroa Marae and Tauranga Bridge Marina monitoring locations, which continued to record occasional exceedances of WHO 10-minute and daily guidelines. The elevated concentrations at these two locations appear to be influenced primarily by SO<sub>2</sub> emissions from the adjacent fertiliser works.

It is less clear what the effects of the elevated daily levels of SO<sub>2</sub> would be. A recent systemic review and meta-analysis concluded that rises in short-term SO<sub>2</sub> concentration increases the risk of all-cause mortality and respiratory mortality (Orellano *et al.*, 2021). This relationship was considered to have a high degree of certainty (WHO, 2021). Another recent systemic review and meta-analysis reported short-term exposure to SO<sub>2</sub> increased the risk of asthma-associated emergency room visits and hospital admissions (Zheng *et al.*, 2021). This relationship is considered causal (WHO, 2021).

The evidence suggests that residents and visitors including manuhiri (guests) to Whareroa Marae and kohanga reo, residents (on boats) and visitors to the Tauranga Bridge Marina may have been, and continue to be, adversely affected by SO<sub>2</sub> emissions.

### *Hydrogen Sulphide*

Ambient air quality monitoring at Whareroa Marae and source investigation by BOPRC provides clear evidence that industrial emissions of hydrogen sulphide (H<sub>2</sub>S) regularly exceeded the national guideline set to prevent against offensive odours (BOPRC, 2020). This would reduce the quality of life and impact adversely on the wellbeing of Marae users, including manuhiri, and Whareroa Marae residents.

### *Benzene*

Limited, short-term monitoring of benzene (C<sub>6</sub>H<sub>6</sub>)<sup>2</sup> in industrial locations of the Mount Maunganui Airshed have measured ambient concentrations above a health-based, air quality

---

<sup>2</sup> Whilst benzene is not a PAH, it is also emitted as a by-product of incomplete combustion (like PAHs) and is also a known carcinogen (like PAHs).



criterion for acute exposure to benzene. Workers in these locations may have elevated acute exposure to benzene, however, the data are limited, and therefore no firm conclusions can be drawn.

We note that Whareroa Marae is within 200 metres of a bulk fuel storage facility and within 300 metres of an oil re-refining facility. Residents and visitors, including manuhiri, to Whareroa Marae and kohanga reo may have been exposed to elevated concentrations of benzene over acute and/or chronic time frames. Benzene exposure may disproportionately impact infants and children (OEHHA, 2014). However, in the absence of any data no firm conclusions can be drawn.

Benzene is a known carcinogen, however, there are no long-term air quality monitoring data for benzene in or around the Mount Maunganui Airshed. This is a significant data gap. No conclusions can be drawn about either worker or residential chronic exposure to benzene.

### *Odour*

Odour is a well-established issue in Mount Maunganui, with more than 500 complaints to the regional council each year.

Based on the frequency of complaints it is apparent that offensive and objectionable odours are reducing the quality of life and adversely impacting on the wellbeing of residents in and around the Mount Maunganui Airshed.

Recent literature suggests that industrial odours are often associated with adverse health impacts in surrounding communities (Government of Alberta, 2017, Guadalupe-Fernandez *et al.*, 2021). Some chemicals, such as benzene, can be harmful even when present below their respective odour thresholds. This suggests that, in addition to negative impacts on wellbeing, odorous emissions may also be adversely impacting residents' health.

### *Data Gaps*

The assessment has highlighted a lack of data for some pollutants of potential concern. Specifically, there is a dearth of ambient air quality monitoring for PAHs and benzene. This has constrained our ability to assess potential cancer risks.

# 1. INTRODUCTION

---

In May 2022, Toi Te Ora Public Health requested the following analysis and advice: :

1. Provide an estimate of health impacts on the population in the Mount Maunganui area from exposure to air contaminants of public health concern where concentrations are known or can reasonably be estimated e.g., particulates (PM<sub>10</sub>, PM<sub>2.5</sub>), sulphur dioxide (SO<sub>2</sub>), methyl bromide (CH<sub>3</sub>Br), hydrogen sulphide (H<sub>2</sub>S). Depending on readily available data possible health impacts for consideration may include:
  - a. Number of excess deaths
  - b. Restricted activity days
  - c. Additional hospital admissions
  - d. Extra presentations for health care
2. Where applicable provide an estimate in the differences in health impacts between compliance with the National Environmental Standards (NES) for Air Quality and meeting relevant health guideline levels.

Toi Te Ora Public Health intends that the information be used to:

- a. Inform polluters, regulatory agencies, and the affected community on the potential scale of adverse health outcomes from current levels of air contaminants; and
- b. Inform parties what reductions in health impacts might be achieved by lowering discharge targets, with the aim of getting the relevant industries and regulators to work together to reduce discharges.

Further to a meeting on 28 June 2022, and following the publication of the third Health and Air Pollution in New Zealand (HAPINZ 3.0) study (Kuschel *et al.*, 2022), Toi Te Ora advised in an email on 18 July 2022 that:

- Nitrogen dioxide (NO<sub>2</sub>), PM<sub>2.5</sub>, SO<sub>2</sub>, H<sub>2</sub>S and polycyclic aromatic hydrocarbons (PAHs) were to be included in the assessment;
- Odour impacts on well-being to be included on a qualitative basis; but
- Methyl bromide was not to be included.

## 1.1 BACKGROUND

Whareroa Marae and the adjoining community are situated on the shore of Tauranga Harbour. The Whareroa Marae is a traditional pa site and key marae for Ngai Tukairangi and Ngāti Kuku hapū of the Ngāi Te Rangi Iwi. Whareroa Pā has been present for around 150 years, making it one of the oldest kāinga (settlements) in the area.

The Mount Maunganui residential area is low lying and represents a relatively constrained geographic area situated between the Tauranga Harbour and the coast. It is a popular residential area but also home to the largest port in New Zealand and a significant industrial estate, generally contained within the Mount Maunganui Airshed – refer **Figure 1**. Whareroa Marae, and some residential areas, pre-date the establishment of the Mount Maunganui industrial estate in the 1940s.



**Figure 2** shows activities sensitive to air pollution in and around the Mount Maunganui Airshed. Of note, there are currently five childcare facilities inside the Mount Maunganui Airshed, including a kohanga reo at Whareroa Marae.

Key industrial activities in the Mount Maunganui Airshed with potentially significant discharges to air currently include:<sup>3</sup>

- Port of Tauranga Ltd (shipping, container, fuel, bulk materials and log loading and unloading)
- Ballance Agri-Nutrients Ltd (sulphuric acid & superphosphate fertiliser manufacture)
- Lawter, Hexion (chemical & resin manufacture)
- Multiple bulk fuel storage and handling facilities
- Multiple bulk materials manufacture, transport, storage & handling (aggregate, cement, coal, fertiliser, stock feed)
- Multiple log transport, storage and handling areas
- Three asphalt plants
- Waste Management Ltd (waste oil refining)

In 2015, Bay of Plenty Regional Council (BOPRC) installed ambient air quality and meteorological monitoring instruments at Whareroa Marae as shown in **Figure 3**. The monitoring revealed significant exceedances of health-based air quality criteria.

In late 2018, BOPRC commenced ambient air quality monitoring at additional locations in the Mount Maunganui Airshed (refer **Figure 3**).

---

<sup>3</sup> Compost and pet food manufacturing activities have recently exited the Mount Maunganui Airshed.

FIGURE 1: Mount Maunganui Airshed (red outline) and adjacent residential areas (orange outline approx. 2013 census area units) [Source: Google Earth, 2 Aug 19]





**FIGURE 2: Early childhood education centres in (yellow stars) and around (red boxes) the Mount Maunganui Airshed (black outline) [Source: Toi Te Ora Public Health]**




FIGURE 3: Locations of Bay of Plenty Regional Council ambient air quality monitoring for PM<sub>10</sub> and SO<sub>2</sub> in the Mount Maunganui Airshed\* [Source: BOPRC]



\*Totara St also measures PM<sub>2.5</sub>, De Havilland Way measures only PM<sub>10</sub>

## 1.2 PRINCIPLES OF ASSESSMENT

This assessment was prepared in accordance with the World Health Organisation (WHO) recommendations for good practice air pollution health risk assessments (WHO, 2014) as follows:

- *Health effects assessment should address an area of uncertainty and an unmet need for information (particularly with respect to social costs).*

The assessment was prepared in response to questions from the local community, industry and Bay of Plenty Regional Council about potential health impacts of air pollution on residents.

- *The assessment reflects the core WHO value of the “right to health”.*

The assessment includes odour, and other pollutants known to be emitted but not monitored in the Mount Manganui Airshed. It considers not only health but also potential impacts on wellbeing.

- *The process of undertaking an assessment is explicit and transparent so the end user can see how health impacts and social costs were selected and calculated.*

Section 2 outlines the methodology for the assessment approach with all references provided.

- *The process of undertaking an assessment is multidisciplinary and includes all relevant expertise and perspectives, including input from stakeholders.*

The quantitative assessment relies on the national health effects model (Sridhar *et al.* 2022) prepared for the HAPNZ Study 3.0 (Kuschel *et al.* 2022). The HAPINZ 3.0 study was prepared by a multidisciplinary team for a steering group comprising officials from the Ministry for the Environment, Ministry of Transport, Waka Kotahi, Ministry of Health and regional councils who all had input to the design and outcomes. The HAPINZ 3.0 study was extensively peer reviewed with strong input from stakeholders.

We understand this assessment will be provided to the community, industry and BOPRC for consideration of wider impacts.

- *The evidence used to develop the assessment is publicly available.*

Cost data were sourced from the Ministry of Transport (refer section 3.2.2).

Exposure response functions developed for New Zealand (Hales *et al.*, 2021) were used for quantitative risk assessment (refer section 3.2.4).

Population data were provided by NZStats (refer section 3.2.5).

Health statistics were provided by the Ministry of Health (refer section 3.2.6).

Air quality and meteorological data were sourced from the BOPRC (refer section 4).

- *Assessment outputs (in the form of exposure and effects models) can be implemented in, and adapted to, local settings and contexts.*

The quantified risk assessment is an adaptation of an existing national tool (HAPINZ 3.0) for the Mount Maunganui setting. Health incidence, air quality and population data are specific to each census area unit.

- *Assessment communication products should be tailored to the general public.*

This document is a technical report outlining, in a transparent manner, the approach and findings. Given the highly technical nature of the material it may not be suitable for the general public, and it is likely that additional material (e.g., summary graphics) will need to be prepared for a lay audience.



## 2. HEALTH EFFECTS OF AIR CONTAMINANTS

---

WHO has identified air quality as the world's largest environmental health risk and among the largest global health risks – comparable with 'traditional' health risks such as smoking, high cholesterol, and obesity. The WHO estimates that indoor and outdoor air pollution exposure currently kills seven million people worldwide every year due to cardiovascular diseases, such as strokes and ischaemic heart disease, as well as respiratory diseases including acute respiratory infections, chronic obstructive pulmonary diseases and lung cancer (WHO, 2021).

The impacts of air pollution are assessed through short-term (acute) or long-term (chronic) exposure. Short-term exposures cover minutes, hours, or days. Long-term exposures are usually over months or years. The major impacts of air pollution occur due to chronic exposure (WHO, 2021).

Accordingly, the quantified assessment of health effects of air pollutants in this document focus on chronic (long-term) exposure.

### 2.1 PARTICULATE MATTER (PM<sub>10</sub> AND PM<sub>2.5</sub>)

Particulate Matter (PM) is a collective term for solid and liquid particles suspended in the air and small enough to be inhaled. The major components of PM are sulphate, nitrates, ammonia, sodium chloride, black carbon, mineral dust and water.

PM is classified by particle size defined through aerodynamic diameter:

- **PM<sub>10</sub>** – particulate less than 10 microns; known as coarse particulate
- **PM<sub>2.5</sub>** – particulate less than 2.5 microns; known as fine particulate.
- **PM<sub>1</sub>** – particulate less than 1 micron; known as ultrafine particulate.

In general, PM<sub>2.5</sub> and smaller tends to be more closely associated with anthropogenic activities, whereas PM<sub>10-2.5</sub> can have a substantial natural source component. The main sources of PM in New Zealand are home heating, industry, agricultural practices, road dust and sea salt. The main anthropogenic (human caused) sources of PM in New Zealand are domestic fires, industry and motor vehicles.

Different sizes of PM can result in different health effects. This is because they deposit in different parts of the respiratory tract, they have diverse sources, and they can interact through different biological mechanisms (WHO, 2013). In general, the smaller a particle is, the farther into the respiratory tract it can penetrate to interact and cause adverse health effects.

There is scientific consensus that exposure to particulate pollution causes predominantly respiratory and cardiovascular effects, ranging from subclinical functional changes (e.g. reduced lung function) to symptoms (increased cough, exacerbated asthma) and impaired activities (e.g. school or work absenteeism) through to doctors' or emergency room visits, hospital admissions and death (WHO, 2006). The effects, in terms of escalating severity, are described as increased visits to doctors for many individuals, hospital admission for some individuals and death for a few individuals. People with pre-existing heart or lung disease, young children, and the elderly, are most likely to suffer adverse health effects. The exposure-response relationship is essentially linear and there is no 'safe' threshold; adverse health effects are observed at all measured levels (US EPA 2020; WHO 2013, WHO, 2021).

In 2013, the International Agency for Research on Cancer (IARC) classified particulate matter (as a component of outdoor pollution) as carcinogenic based on an increased risk of lung cancer (IARC, 2013). Additional research further indicates particulate matter is associated with atherosclerosis, adverse birth outcomes, childhood respiratory disease (WHO, 2013) as well as Alzheimer’s disease and other neurological endpoints, cognitive impairment, diabetes, systemic inflammation and aging (WHO 2016b).

More recently, WHO has concluded that chronic exposure to PM is causal, or likely to be causal, for (WHO, 2021):

- All-cause mortality
- Cardiovascular mortality (all, cerebrovascular, ischaemic heart disease)
- Respiratory mortality (any, chronic obstructive pulmonary disease, acute lower respiratory infections)
- Lung cancer

**Table 1** presents New Zealand and WHO air quality guidelines and standards for particulate matter.

**TABLE 1: New Zealand and WHO air quality guidelines and standards for particulate matter**

Time Average / Jurisdiction (Year)	Guideline / Standard ( $\mu\text{g}/\text{m}^3$ )	Permitted Exceedances per Year
<b>PM<sub>2.5</sub></b>		
24-hours		
New Zealand (2002)*	25	–
WHO (2021)	15	3 – 4
Annual		
WHO (2021)	5	–
<b>PM<sub>10</sub></b>		
24-hours		
New Zealand (2004)	50	1
WHO (2021)	45	3 – 4
Annual		
New Zealand (2002)	20	–
WHO (2021)	15	–

\*Reporting guideline only

## 2.2 NITROGEN DIOXIDE (NO<sub>2</sub>)

Nitrogen dioxide (NO<sub>2</sub>) is a reddish brown coloured acidic gas with a characteristic pungent odour. The main sources of NO<sub>2</sub> worldwide are combustion processes such as motor vehicles, domestic heating, industrial combustion sources, electricity generation, shipping and construction machinery. NO<sub>2</sub> is both a primary and secondary pollutant i.e., it is both emitted and forms downwind from other pollutants (including PM<sub>2.5</sub>).

Nitrogen dioxide is the main source of nitrate aerosols, which form an important fraction of PM<sub>2.5</sub> and, in the presence of sunlight, ozone. It is also a major component of brown haze. In

New Zealand the main source of nitrogen dioxide is motor vehicles. In Mount Maunganui, ships and industry are also likely to be significant sources of nitrogen dioxide.

Long-term exposure to NO<sub>2</sub> increases the risk of premature death (mortality) and respiratory illnesses (morbidity) (WHO, 2021). Epidemiological studies have also shown that symptoms of bronchitis in asthmatic children increase with long-term exposure to NO<sub>2</sub>. Reduced lung function is also linked to measured levels within cities of Europe and North America (WHO, 2006). Recent evidence also suggests exposure may increase the risk of premature death and trigger heart attacks (USEPA, 2016).

Short-term exposure to high concentrations of nitrogen dioxide (NO<sub>2</sub>) causes significant inflammation of the airways and respiratory problems and can also trigger asthma attacks (WHO, 2021).

**Table 2** presents New Zealand and WHO air quality guidelines and standards for NO<sub>2</sub>.

**TABLE 2: New Zealand and WHO quality guidelines and standards for NO<sub>2</sub>**

Time Average / Jurisdiction (Year)	NO <sub>2</sub> Guideline / Standard (µg/m <sup>3</sup> )	Permitted Exceedances per Year
1-hour		
New Zealand (2004)	200	9
WHO (2000)	200	–
24-hours		
New Zealand (2002)	100	–
WHO (2021)	40	3 – 4
Annual		
WHO (2021)	10	–

### 2.3 SULPHUR DIOXIDE (SO<sub>2</sub>)

Sulphur dioxide is a colourless gas that is readily soluble in water. It has a characteristic pungent smell and an odour threshold of 870 micrograms per cubic metre (µg/m<sup>3</sup>) or 0.33 parts per million (ppm) (AIHA, 2013). The odour threshold is the level at which 50% of the population can just detect the odour.

Sulphur dioxide is both a primary and secondary pollutant, it is a precursor for the formation of PM<sub>2.5</sub>. Sulphur dioxide arises naturally from volcanic sources, with White Island being New Zealand's largest source of sulphur dioxide (PAE, 2009). Away from industrial and volcanic sources, background levels of sulphur dioxide in New Zealand are typically very low, less than 5 µg/m<sup>3</sup> as a one-hour average.

In New Zealand the major anthropogenic sources are industrial processes (aluminium manufacture, fertiliser manufacturing, chemicals manufacture) and the combustion of fossil fuels that contain sulphur. The main sources of sulphur dioxide in Mount Maunganui are shipping (745 tonnes/year) and industry (232 tonnes/year) (Wilton & Iseli, 2019). It should be noted that the inventory estimate of shipping SO<sub>2</sub> emissions was for the year 2018 and predates the implementation of Annex VI of MARPOL on 1 January 2020.<sup>4</sup>

<sup>4</sup> This requires ships to burn low sulphur fuel or implement abatement technology to mitigate emissions of SO<sub>2</sub>.

Sulphur dioxide can cause respiratory problems, such as bronchitis, and it can irritate the nose, throat and lungs. This is because inhaled sulphur dioxide readily reacts with the moisture of mucous membranes to form sulphurous acid (which is a severe irritant). It may cause coughing, wheezing, phlegm and asthma attacks (MfE, 2011).

Studies have shown that asthmatics and people with lung disease are particularly sensitive to sulphur dioxide. Children may also be more sensitive to the effects of sulphur dioxide due to their relatively higher respiration rate and smaller body mass.

In 2021, WHO published two systematic reviews and meta-analyses on the effects of short-term exposure to ambient sulphur dioxide (SO<sub>2</sub>) on:

- all-cause and respiratory mortality (Orellano *et al.*,2021); and
- emergency room visits and hospitalisations for asthma (Zheng *et al.*,2021).

Orellano *et al.*,2021 found that **short term increases in SO<sub>2</sub> increased the risk of all-cause mortality (daily SO<sub>2</sub>) and respiratory mortality (1-hour SO<sub>2</sub>) with a high certainty of evidence.** In general, concentration response functions showed linear behaviour with no thresholds. Orellano *et al.*,2021 considered the epidemiological evidence supports a causal relationship.

Zheng *et al.*,2021 found that short term increases in SO<sub>2</sub> correlate with increased risk of asthma-associated emergency room visits and hospital admissions. Children and to a lesser extent the elderly are more susceptible. The positive correlation between daily SO<sub>2</sub> and asthma-associated emergency room visits and hospital admissions was judged as having a moderate certainty of evidence and warrants further investigation. SO<sub>2</sub> was not found to have a daily threshold of effects.

The speed with which people show health effects from exposure to SO<sub>2</sub> necessitates a focus on acute exposure. National environmental standards and ambient air quality guidelines for SO<sub>2</sub> are presented in **Table 3**.

**TABLE 3: New Zealand and WHO short-term air quality guidelines and standards for SO<sub>2</sub>**

Time Average / Jurisdiction (Year)	SO <sub>2</sub> Guideline / Standard (µg/m <sup>3</sup> )	Permitted Exceedances per Year
10-minutes WHO (2000)	500	0
1-hour New Zealand (2004)	350 570	9 0
24-hours New Zealand (2002) WHO (2021)	120 40	– 3 – 4

## 2.4 HYDROGEN SULPHIDE (H<sub>2</sub>S)

Hydrogen sulphide (**H<sub>2</sub>S**) is a flammable, colourless gas with the characteristic odour of rotten eggs. H<sub>2</sub>S is odorous at extremely low concentrations (0.06 µg/m<sup>3</sup>) (AIHA, 2019). H<sub>2</sub>S gas is found naturally in geothermal areas and is also emitted from volcanoes, undersea vents, swamps and stagnant bodies of water. H<sub>2</sub>S is also emitted from industrial processes such as oil refining, pulp and paper manufacture, tanneries and wastewater treatment.

H<sub>2</sub>S is heavier than air and can build up in confined or low-lying, still spaces. Sanitation workers are most at risk from exposure to H<sub>2</sub>S when working in or near sewers, septic tanks and sump holes, however, geothermal areas, such as Rotorua, also pose risks to the general public. There have been 11 deaths linked with H<sub>2</sub>S exposure in geothermal spa pools and confined spaces in New Zealand between 1946 and 2008.<sup>5</sup>

H<sub>2</sub>S is a gas that affects the nervous system and cardiovascular system leading to a range of symptoms such as nausea, headache and dizziness. The physiological response is rapid, even one or two breaths at high concentrations (> 75,000 – 150,000 µg/m<sup>3</sup>) can cause ‘knock down’. Single exposures to very high concentrations may rapidly cause breathing difficulties and death (PHE 2009).

WHO, 2003 notes:

*Health effects that have been reported as being associated with odour exposures include nausea, headaches, retching, difficulty breathing, frustration, annoyance, depression, stress, tearfulness, reduced appetite, and being woken in the night. There are also social effects such as reduced enjoyment of the outdoors and embarrassment in front of visitors. All of these contribute to a reduced quality of life for the individuals who are exposed. People can also develop physiological effects from odour even when their exposure is much lower than that normally associated with the reported physical health effects. This effect is sometimes termed ‘odour worry’ and is due to the perception that if there is a smell it must be doing physical harm.*

The New Zealand guideline for H<sub>2</sub>S is 7 µg/m<sup>3</sup> as a 1-hr average (MfE, 2002). This was set to prevent odour annoyance and the resulting impacts on well-being, rather than specific health effects.

For context, the California Office of Environmental Hazard and Health Assessment (OEHHA) has established a reference exposure level (REL) of 42 µg/m<sup>3</sup> as a 1-hour average that it is “protective against mild adverse effects (headaches and nausea)” noting that at this concentration around 40% of the general population would find H<sub>2</sub>S to be objectionable (OEHHA 1999).<sup>6</sup>

**Table 4** presents health effects of H<sub>2</sub>S at varying concentrations.

---

<sup>5</sup> Rotorua Daily Post (2008). *The Death Toll*. 13 June 2008.

<sup>6</sup> Appendix D2. Acute RELs and toxicity summaries using the previous version of the Hot Spots Risk Assessment guidelines.

**TABLE 4: Acute health effects of H<sub>2</sub>S at varying concentrations** [Source: ESR 2022]

Concentration (µg/m <sup>3</sup> )	Symptoms/Effects
0.2 – 0.5	Typical background concentrations
0.06 – 2,130*	Odour threshold (rotten egg smell is noticeable)
3,000 – 7,500	<ul style="list-style-type: none"> <li>- Odour becomes more offensive</li> <li>- Prolonged exposure may cause nausea, tearing of the eyes, headaches or loss of sleep.</li> <li>- Airway problems (bronchial constriction) in some asthma patients</li> </ul>
30,000	<ul style="list-style-type: none"> <li>- Possible fatigue</li> <li>- Irritability</li> <li>- Above 45,000 µg/m<sup>3</sup> odour described as sweet or sickeningly sweet</li> <li>- Loss of appetite</li> <li>- Poor memory</li> <li>- Headache</li> <li>- Dizziness</li> </ul>
75,000 – 150,000	<ul style="list-style-type: none"> <li>- Slight conjunctivitis (“gas eye” or pink eye) and respiratory tract irritation after 1 hour</li> <li>- May cause digestive upset and loss of appetite</li> </ul>
150,000	<ul style="list-style-type: none"> <li>- Coughing, eye irritation, loss of smell after 2-15 minutes (olfactory fatigue)</li> <li>- Altered breathing, drowsiness after 15-30 minutes</li> <li>- Throat irritation after 1 hour</li> <li>- Gradual increase in severity of symptoms over several hours</li> <li>- Death may occur after 48 hours</li> </ul>
150,000 – 230,000	<b>Loss of smell (olfactory fatigue or paralysis)</b>
300,000 – 450,000	<ul style="list-style-type: none"> <li>- Marked conjunctivitis and respiratory tract irritation after 1 hour</li> <li>- Pulmonary oedema may occur from prolonged exposure</li> </ul>
750,000 – 1,000,000	<ul style="list-style-type: none"> <li>- Staggering, collapse in 5 minutes</li> <li>- Serious damage to the eyes in 30 minutes</li> <li>- Death after 30-60 minutes</li> </ul>
1,000,000 – 1,500,000	<ul style="list-style-type: none"> <li>- Rapid unconsciousness</li> <li>- “Knockdown” or immediate collapse within 1 to 2 breaths</li> <li>- Breathing stops</li> <li>- Death within minutes</li> </ul>
1,500,000 – 3,000,000	Nearly instant death

\*AIHA, 2019

## 2.5 BENZENE

Benzene (C<sub>6</sub>H<sub>6</sub>) is a colourless, clear liquid with a boiling point of 80°C (MfE, 2002).

Motor vehicles and household fires are significant sources of benzene in New Zealand’s air. There are also some industrial activities that use and discharge benzene. Motor vehicle exhaust emissions of benzene are thought to derive partly from unburnt benzene in the fuel, and partly from the dealkylation of other aromatic hydrocarbons (MfE, 2002). In the Mount Maunganui Airshed, petroleum re-refining, tank filling and ‘breathing’ losses from petroleum storage are likely sources of benzene. ‘Breathing’ losses refer to vapours in the headspace of



the storage tank discharging to atmosphere due to changes in the tank temperature and pressure throughout the day and throughout the year.

Benzene is a known human carcinogen (IARC Group 1; leukaemia) and is known to have haemotoxic, genotoxic & mutagenic effects. The New Zealand guideline for benzene is 1 part per billion (ppb) as an annual average (MfE, 2002).<sup>7</sup> This was set on a precautionary basis for a carcinogen in 2002.

With respect to non-cancer effects, acute, high inhalation exposure to benzene may lead to eye, nose, and throat irritation and central nervous system depression in humans. Prolonged or repeated exposures have been associated with both blood cell proliferation and reduction in blood cell numbers due to bone marrow suppression, including peripheral lymphocytopenia, pancytopenia, and aplastic anaemia (OEHHA 2014).

OEHHA determined there is valid concern that benzene exposure may disproportionately impact infants and children. This is based on the wide-spread exposure to benzene, the documented toxicokinetic variability in benzene metabolism, the transplacental genotoxicity and developmental toxicity of benzene, the documented increased sensitivity of early in life exposure to benzene carcinogenicity in animals, as well as the dynamic haematopoiesis that occurs during human development (OEHHA, 2014).

### 2.5.1 Chronic Benzene Criteria

For known carcinogens good practice in New Zealand is to quantitatively estimate risk using factors for carcinogens provided by the United States Environment Protection Agency (US EPA) Integrated Risk Information System (IRIS) database. The IRIS database has been developed from toxicological data using a rigorous and transparent methodology with an ongoing process of review and revision. The recommended approach for air quality assessments in New Zealand is to adopt an acceptable environmental risk for exposure by residential receptors to individual environmental pollutants of 1 in 1,000,000 (i.e., one in a million).<sup>8</sup>

The US EPA inhalation unit risk factor for benzene is  $2.2 \times 10^{-6}$  per  $\mu\text{g}/\text{m}^3$ , set to protect against leukaemia.<sup>9</sup>

### 2.5.2 Acute Benzene Criteria

New Zealand has no acute (short-term) air quality guideline for benzene. In the absence of any local guidance, good practice is to refer to criteria established using transparent derivation from toxicological data, whilst explaining the purpose of the guideline (MfE, 2016).

The OEHHA reference exposure levels (RELs) have been developed from toxicological data using a rigorous and transparent methodology with an ongoing process of review and revision to take into account new information and sensitive subpopulations including infants and children. Acute RELs are concentrations that are not likely to cause adverse effects in a human population, including sensitive subgroups, exposed to that concentration on an intermittent basis for one hour. Acute RELs are intended to protect the individuals who live or work in the vicinity of emissions of these substances. The focus of acute RELs is generally a one-hour exposure for *non-cancer health impacts*.

---

<sup>7</sup>  $3.6 \mu\text{g}/\text{m}^3$  when converted at  $0^\circ\text{C}$ , 101.3 kPa (MfE, 2009)

<sup>8</sup> MfE, 2016. At section 4.5 (page 54).

<sup>9</sup> [https://iris.epa.gov/ChemicalLanding/&substance\\_nmbr=276](https://iris.epa.gov/ChemicalLanding/&substance_nmbr=276). Accessed 13 Dec 2022.



The OEHHA has set a 1-hour REL for benzene of 8 parts per billion (ppb, OEHHA, 2014).<sup>10</sup> This was set to protect against developmental effects.

## 2.6 ODOUR

Current good practice guidance states (MfE, 2016b):

*“People have reported effects of odour that include nausea, headaches, retching, difficulty breathing, frustration, annoyance, depression, stress, tearfulness, reduced appetite, being woken in the night, and embarrassment in front of visitors. All of these contribute to a reduced quality of life for the individuals who are exposed to the odour and underscore the importance of taking odour complaints seriously.”*

Traditionally odour has been considered an amenity issue because of the lack of any relationship with toxicological effects. However, there is a nascent body of epidemiology associating adverse odours from industrial sources with adverse health effects.

A review of more than 50 studies published between 1975 and 2013 found that residents of communities located near odour emitting facilities were found to report a higher number of health symptoms compared to residents of control communities (Government of Alberta, 2017). Reported outcomes included respiratory symptoms, nausea, congestion, eye irritation, headache, dizziness, sleep problems, and diarrhoea. These symptoms were observed in response to odours from a range of sources including petroleum refineries, livestock operations, hazardous waste sites, municipal landfills, and industrial plants.

More recently, a systematic review and meta-analysis of 30 odour studies found a statistically significant association between populations exposed to odour pollution and adverse health effects such as headache and cough/phlegm (Guadalupe-Fernandez *et al.*, 2021), noting:

*“Meta analysis results showed that residential odour exposure was associated to an increased risk of headache and cough/phlegm, and to a borderline risk of nausea and vomiting. We found suggestive associations for the other outcomes investigated (e.g., asthma, mucus irritation, mood states) but evidence is sparse.*

*The associations with headache, cough/phlegm and nausea/vomiting have a biological plausibility. Unpleasant odours are able to modulate autonomic system responses, such as vagal nerve inducing nausea or vomiting [5]. Another mechanism involves stress, consequent to environmental worry [18], and stress-related psychosomatic reactions such as chronic muscular tension, headaches, sleep disturbance. Chemicals responsible for odour may cause irritation, supporting the higher risk for cough/phlegm. Eye and nose irritation and asthma exacerbations can also be related to this odour-related irritation, but only limited evidence was found in this review.”*

---

<sup>10</sup> 27 µg/m<sup>3</sup> when converted at 0°C, 101.3 kPa (OEHHA, 2014)



### 3. METHODOLOGY


A health hazard can be defined as a source of risk to human health or wellbeing. A health risk assessment is the scientific evaluation of potential adverse health effects resulting from human exposure to a particular hazard (WHO 2016a). In the context of this document, the health hazard of interest is air pollution.

An air pollution health risk assessment (AP-HRA) aims to estimate the risks of past, current or future exposure to air pollution. An AP-HRA may be quantitative or qualitative; it assesses:

- The amount of air pollution present (i.e., pollutant concentrations);
- The amount of contact (exposure) of the selected population; and
- How harmful the pollutant concentrations are for human health.

Figure 4 provides an overview of the AP-HRA process.

FIGURE 4: Overview of an Air Pollution Health Risk Assessment process [Source: WHO 2016a]



While an AP-HRA tends to look into particular hazards and their effects on human health, a health impact assessment (HIA) takes a broader perspective. For example, when considering the impacts of industry in Mount Maunganui on public health, an HIA would look into not only the quantitative and qualitative risks associated with discharges to air, but also issues such as noise and soil and water pollution, as well as the positive economic benefits on the population from the Port of Tauranga and associated business.

This assessment is more akin to an AP-HRA as opposed to a broader HIA considering the wider impacts (i.e., both positive and negative impacts). It seeks to address the policy question:

*“What are the health impacts of air contaminants of public health concern in the Mount Maunganui area?”*

### **3.1 CONTAMINANTS OF PUBLIC HEALTH CONCERN**

Toi Te Ora Public Health specified the following contaminants for assessment:

- Hydrogen sulphide (H<sub>2</sub>S)
- Nitrogen dioxide (NO<sub>2</sub>)
- Particulates (PM<sub>10</sub>, PM<sub>2.5</sub>)
- Polycyclic aromatic hydrocarbons (PAHs)
- Sulphur dioxide (SO<sub>2</sub>)
- Odour

Polycyclic aromatic hydrocarbons (PAHs) are a class of organic compounds composed of multiple aromatic rings. PAHs are produced by incomplete combustion of organic material, for example in ships and vehicles or domestic home heating. PAHs generally have a low degree of acute toxicity to humans following exposures of short duration. Repeated exposure to PAHs over a long(er) period of time, usually in occupational settings, has been associated with increased incidence of lung, skin, and bladder cancers. It is difficult to ascribe observed health effects in epidemiological studies to specific PAHs because most exposures are to PAH mixtures (ATSDR 2009). Any carcinogenic risk associated with PAHs will be a function of exposure intensity (concentration) and exposure duration.

We are not aware of any emissions data or ambient air quality monitoring for PAHs in the Mount Maunganui Airshed. Therefore, PAHs were not able to be included in this assessment.

We are, however, aware that some industries undertake intermittent (screening), speciated, ambient monitoring for volatile organic compounds (VOCs), including benzene, and report this to BOPRC. Benzene is a known carcinogen, and (like PAHs) it is emitted as a product of incomplete combustion from ships and vehicles. Being volatile, benzene is also emitted from bulk fuel storage and transfer. Given the presence of bulk fuel storage at the Port of Tauranga we consider benzene may be a contaminant of public health concern in the Mount Maunganui Airshed. We have therefore, included benzene in this assessment.

We also note that hydrogen fluoride (HF) is discharged to air by the fertiliser works, which is adjacent to Whareroa Marae. Unfortunately, due to monitoring instrument failure, there are insufficient data to assess current levels of hydrogen fluoride. Previous reporting by BOPRC (BOPRC 2020) has shown regular exceedances of the national daily general land use critical

level for protecting ecosystems for hydrogen fluoride ( $2.9 \mu\text{g}/\text{m}^3$ , MfE 2002) at the Whareroa Marae monitoring site.

### 3.2 QUANTIFIED RISK ASSESSMENT: $\text{PM}_{10}$ , $\text{PM}_{2.5}$ AND $\text{NO}_2$

Air pollution health risk assessments typically compare the estimated health effects in a population exposed to a source of concern, with the estimated health effects in the same population without the source of concern. This is known as the counterfactual approach.

The source of concern here is shipping and industrial emissions in the Mount Maunganui Airshed. Therefore, the counterfactual approach would be to compare the estimated health effects of air quality in Mount Maunganui with and without industry or shipping emissions. This is somewhat hypothetical. A slightly more realistic comparison was adopted in this assessment.

The approach in this assessment was to compare the estimated health effects in a population exposed to shipping and industrial emissions (i.e., air quality in Mount Maunganui) with the estimated health effects in a similar population without these emissions (i.e., air quality in Otūmoeti).

A valid criticism of this approach is that Otūmoetai is located close to the Mount Maunganui Airshed (around 5 km) and may similarly be affected by emissions from industrial and shipping activity. We are comfortable that this means the estimates are not conservative and likely underestimate effects.

Modelling was undertaken using the Health and Air Pollution in New Zealand (HAPINZ 3.0) exposure model (Sridhar *et al.*, 2022). The HAPINZ 3.0 exposure model has New Zealand specific exposure response functions for key air pollutants developed by Hales *et al.* (2021).

#### 3.2.1 Calculating the health burden

For each area under assessment, the health impacts are estimated as follows:

$$\text{Health Effects (cases)} = \text{Cases (total)} \times \text{PAF}$$

Where:

**Health effects (cases)** are the number of deaths, hospital admissions or restricted activity days (depending on the health outcome being assessed) due to air pollution.

**Cases (total)** is the total number of health cases (e.g., deaths, hospital admissions) in the area of interest (i.e., health incidence data based on analysis of Ministry of Health mortality and hospitalisations datasets by census area unit).

**PAF** (population attributable fraction) is the estimated percentage of total health cases that are attributable to the air pollution exposure.

The PAF is calculated using the exposure–response function (the relative increase in the health effect for every increment of air pollution (for example, 1.105 for every  $10 \mu\text{g}/\text{m}^3$  of annual average  $\text{PM}_{10}$ )<sup>11</sup> and the exposure (the average pollution concentration in the area of interest - for example, an annual average  $\text{PM}_{10}$  concentration of  $20 \mu\text{g}/\text{m}^3$ ).

---

<sup>11</sup> A relative risk of 1.105 means the risk increases by 10.5% per pollution increment, in this case per  $10 \mu\text{g}/\text{m}^3$  of annual average  $\text{PM}_{10}$ .

This approach estimates the health effects that would be prevented if exposure to the pollutant (e.g., PM<sub>10</sub>) was at the minimum risk level possible, recognising there being no safe threshold for most air pollutants.

### 3.2.2 Estimating social costs

Mortality and morbidity impacts amount to loss of life and loss of quality of life for people exposed to air pollution. These costs can be estimated to arrive at a total cost to society (the *social* cost) resulting from exposure to air pollution. Adverse effects resulting from air pollution place a significant burden on society and health systems (Kuschel et al., 2022).

The social costs of air pollution are then calculated as follows:

$$\text{Social Costs} = \text{Health Effects (cases)} \times \text{Cost per case}$$

In simple terms, the health effects cases estimated as per the previous formula (for example, the number of premature deaths) are combined with published health-cost data to estimate costs.

#### Mortality

We used a value of statistical life (VoSL) approach, whereby the estimated change in the number of premature deaths was multiplied by the current New Zealand VoSL based on transport risk (road safety), as was done in HAPINZ 3 (Kuschel et al., 2022). This assessment utilises a value of statistical life of \$4,527,300 per death (MoT 2020).

The 95% confidence interval estimates of a value of statistical life were \$4,050,742 - \$5,242,137. All costs are in \$NZD 2019.

#### Morbidity

The social cost of hospital admissions were estimated from the financial costs per hospitalisation (based on the average number of bed nights for each type of disease), productivity losses from time off work or school for those hospitalised, family and friends, and recovery costs after discharge from hospital including any long-term disability (quality of life). Full details and assumptions for estimates are provided in Kuschel et al., 2022 and are not repeated here.

The total costs per case with low and high-end estimates were:

- \$36,666 (\$10,809, \$473,294) for cardiac admissions; and
- \$31,748 (\$5,891, \$462,770) for respiratory admissions.

The social cost of restricted activity days (RADs) was \$89 per case, estimated based on lost average income per day. The low and high-end cost estimates for sensitivity analysis were \$49 and \$125 respectively.

Childhood asthma costs included those resulting from general practitioner visits, medication (\$128 per case) and hospitalisations (\$1,822 per case). The low and high-end cost estimates for sensitivity analysis were arbitrarily increased or decreased by 50%. All costs are in \$NZD 2019.

### 3.2.3 Base year

Council commenced monitoring for PM<sub>10</sub> in 2014 in Otūmoetai. To undertake this investigation it was initially proposed to use an extended time average (e.g., a ten-year period ending 2021) for the period of assessment. This would increase the stability of the key exposure metric, ambient air pollutant concentrations. It may, however, underestimate the potential population

exposed (which has increased during this period) and overestimate health impacts (some key incidence statistics have decreased). However, there are insufficient data for the Mount Maunganui Airshed, where monitoring of PM<sub>10</sub> and PM<sub>2.5</sub> only commenced at the end of 2018, to make this feasible. The 2019 – 2021 period was used as the base period for assessment.

Council commenced monitoring for total suspended particulate (TSP) at Totara Street in 2015, with a full year of data available from 2016. Although TSP will include some larger size fractions (up to 100 µm), it is generally taken to represent particulate matter less than 30 µm (i.e., PM<sub>30</sub>). PM<sub>10</sub> is, therefore, a component of TSP. **Figure 5** presents annual average PM<sub>10</sub> concentrations measured in (residential) Otūmoetai (2014 – 2021) for comparison with TSP and PM<sub>10</sub> measured in the (industrial parts of) Mount Maunganui Airshed (2019 – 2021).

**FIGURE 5: Annual total suspended particulate (TSP) and PM<sub>10</sub> measured in Otūmoetai and select Mount Maunganui industrial locations** [Source: BOPRC]




Figure 5 shows there were reductions in annual PM concentrations measured at all industrial locations<sup>12</sup> in the Mount Maunganui Airshed in 2020 and 2021 compared with 2019. The air quality expert consensus view<sup>13</sup> is that these reductions were due to a combination of improved emissions control by the Port and reductions in shipping emissions required under international law (specifically Annex VI of MARPOL).<sup>14</sup> However, these reductions may also reflect changes due to the COVID-19 pandemic which, through stay-at-home restrictions, may have reduced long-term concentrations of key pollutants in and around the Mount Maunganui Airshed.

Irrespective, using data averaged over the three years ending 2021 may significantly underestimate public exposure in previous and future years. We have therefore, selected the year 2019 as the base year for assessment purposes. This is the most recent year for which there is comprehensive ambient air quality monitoring data, but which excludes step changes

<sup>12</sup> Specifically, reductions in annual average PM<sub>10</sub> of 11% (Rata Street), 24% (Rail Yard South) and 15% (Totara Street).

<sup>13</sup> Joint Witness Statement of Air Quality Experts Dr Emily Wilton, Mx Lou Wickham, Ms Jenny Simpson, Mr Paul Baynham, Mr Peter Stacey & Mr Andrew Curtis dated 27 May 2021 for the Environment Court ENV-2019-AKL-000065 and ENV-2019-AKL-000073. At [Q15].

<sup>14</sup> In force from 1 January 2020.

in emissions due to mitigation, regulatory changes and/or any changes due to activity restrictions from COVID-19. Where available, population and health data have been averaged over 2018-2019.

### 3.2.4 Exposure response functions

The mortality and morbidity dose response functions for PM<sub>2.5</sub> and NO<sub>2</sub> are from Hales *et al.*, 2021 which was a New Zealand (national) cohort study.<sup>15</sup> This means that these dose response functions provide a robust estimate of health effects of exposure to air pollution across New Zealand using PM<sub>2.5</sub> and NO<sub>2</sub>, together, as indicators of exposure to air pollution. These were (all Hales *et al.*, 2021 except where noted):

#### PM<sub>2.5</sub>

- Premature mortality risk (per 10 µg/m<sup>3</sup>) for all adults (30+years) associated with annual PM<sub>2.5</sub> exposure 1.105 (95% CI 1.065 – 1.145)
- Cardiovascular hospitalisation risk (per 10 µg/m<sup>3</sup>) for all ages associated with annual PM<sub>2.5</sub> exposure 1.115 (95% CI 1.084 – 1.146)
- Respiratory hospitalisation risk (per 10 µg/m<sup>3</sup>) for all ages associated with annual PM<sub>2.5</sub> exposure 1.070 (95% CI 1.021 – 1.122)
- Restricted activity days risk (per 10 µg/m<sup>3</sup>) for all ages associated with annual PM<sub>2.5</sub> exposure 0.9 (lower/upper bounds 0.5 – 1.7, Ostro, 1987)

#### NO<sub>2</sub>

- Premature mortality risk (per 10 µg/m<sup>3</sup>) for all adults (30+years) associated with annual NO<sub>2</sub> exposure 1.097 (95% CI 1.074 – 1.120)
- Cardiovascular hospitalisation risk (per 10 µg/m<sup>3</sup>) for all ages associated with annual NO<sub>2</sub> exposure 1.047 (95% CI 1.031 – 1.064)
- Respiratory hospitalisation risk (per 10 µg/m<sup>3</sup>) for all ages associated with annual NO<sub>2</sub> exposure 1.130 (95% CI 1.102 – 1.159)
- Asthma prevalence risk (per 4 µg/m<sup>3</sup>) for 0 – 18-year-olds associated with annual NO<sub>2</sub> exposure 1.05 (95% CI 1.02 – 1.07, Khreis *et al.*, 2017)<sup>16</sup>

The premature mortality dose response function for PM<sub>10</sub> is also from Hales *et al.*, 2021. This means that the dose response function provides a robust estimate of health effects of exposure to air pollution across New Zealand using PM<sub>10</sub> as an indicator of exposure to air pollution. This was:

- Premature mortality risk (per 10 µg/m<sup>3</sup>) for all adults (30+years) associated with annual PM<sub>10</sub> exposure 1.111 (95% CI 1.089 – 1.133)

NB: The exposure response function for PM<sub>10</sub> is from a single pollutant model and is separate to the exposure response functions developed by Hales *et al.*, 2021 for PM<sub>2.5</sub> and NO<sub>2</sub> from a two-pollutant model. **This means that the estimate of effects associated with PM<sub>10</sub> is not**

---

<sup>15</sup> With two additions; restricted activity days (Ostro, 1987) and prevalence of asthma (Khreis *et al.*, 2017)

<sup>16</sup> Toi Te Ora requested an assessment of additional presentations for health care, which was not estimated by Hales *et al.*, 2021. However, asthma prevalence in children (under 18-year-olds) associated with long-term NO<sub>2</sub> was assessed and is included here.



**additive to the estimate of effects associated with PM<sub>2.5</sub> and NO<sub>2</sub>, rather it is a separate estimate.**

The specific mix and composition of air pollutants in and around Mount Maunganui is likely to be quite different to typical urban air pollution in New Zealand. This is because typical urban air pollution in New Zealand is influenced by traffic (all year round) and domestic solid fuel burning during winter (only). In Mount Maunganui, however, long-term levels of:

- PM<sub>10</sub> are likely to be dominated by emissions from the Port and industrial activities.
- PM<sub>2.5</sub> and NO<sub>2</sub> are likely to be dominated by emissions from ships and transport, with some influence from industry.

This means that the selected exposure response functions may not accurately estimate the effects of air pollution in and around the Mount Maunganui areas.

It is uncertain whether a Mount Maunganui dose response function would increase or decrease the estimates. However, given the relatively small population exposed (<50,000), it is not feasible to develop a specific dose response function for these pollutants in this locale, and the New Zealand specific functions are the best available estimates.

### **3.2.5 Population assessed**

**Figure 1** presents the NZ Stats 2013 census area units<sup>17</sup> (orange outline) assessed and the Mount Maunganui Airshed (red outline).<sup>18</sup> We selected the five census area units in and around Mount Maunganui (light blue shaded areas) that include the Mount Maunganui Airshed for assessment. These are:

- Mount Maunganui North
- Omanu
- Tauranga City-Marinas
- Arataki
- Sulphur Point

We also assessed the following four (orange NZ Stats 2013) census area units as representative of the Otūmoetai residential area for comparison:

- Matua
- Otūmoetai North
- Otūmoetai South
- Bellevue

The HAPINZ national exposure model has population statistics by census area unit for a base year of 2016. This was updated using StatsNZ usual resident population data averaged for the period 2018 – 2019 (Metcalf and Kuschel, pending). The usual resident population does not include the working population present in Mount Maunganui each day of the working week.

---

<sup>17</sup> Health statistics are disaggregated to 2013 census area units and are not available by 2018 (census) statistical area units.


<sup>18</sup> Approximate only for illustrative purposes

2018 census data suggests this is around an additional 10,000 people<sup>19</sup> which may be compared with the usual resident population of the Mount Maunganui and Otūmoetai areas under assessment (around 35,000 in total) - refer **Figure 6**.

Whilst the working population is only present for around 40 hours per week, they are likely to be exposed to higher daily and annual concentrations of some pollutants than residents as they are physically closer to the primary sources (transport and industry). However, some of the working population may also be wearing personal protective equipment at least some of the time.

The exclusion of the exposed working population is likely to underestimate potential exposure and therefore, likely underestimate associated effects.

**FIGURE 6: Where we live versus where we work (using 2018 census data) (Source NZ Stats (2020))<sup>20</sup>**



<sup>19</sup> 2018 Census employed statistical areas (SA) and did not employ census area units (CAU) so a direct comparison cannot be made. Estimate of around 10,000 comprises SA2 (Mount Maunganui Central), SA1 – 7013954 and SA1 – 7013955 main means of travel to work (2)(4)(7), by workplace address (12) for the employed census usually resident population count aged 15 years and over (5)

<sup>20</sup> <https://storymaps.arcgis.com/stories/6f8b5f981ad34f11bedaf1725e9cb698>



### 3.2.6 Health statistics


Health statistics by census area unit were provided in HAPINZ 3 and sourced, in turn, from the New Zealand Mortality Collection data (MoH 2021a) and the National Minimum Dataset of publicly funded hospital discharges (MoH 2021b).

The health incidence baseline data in the national health effects model is for a base year 2016 (Sridhar *et al.*, 2022).<sup>21</sup> We used updated mortality and morbidity data for the two-year period 2018 – 2019 (Metcalf and Kuschel, pending). It should be noted that our assessment focusses (only) on the relevant census area units with this updated data.

Historical mortality statistics (MoH, 2021c) reveal a slow but steady decline in mortality rates as shown in **Figure 7**. This means that using 2019 as a base year is not likely to overestimate effects.

It should be noted that the impacts of COVID-19 are excluded from this assessment, not least because mortality statistics for 2020 and 2021 will not be available until 2023 and 2024 respectively.

**FIGURE 7: National mortality rates 2009-2018** [Source: New Zealand Mortality Collection Records 1996-2018]



### 3.3 QUALIFIED RISK ASSESSMENT: SO<sub>2</sub>, H<sub>2</sub>S, BENZENE AND ODOUR

The approach taken was to review the available ambient air quality monitoring data for comparison with health-based standards and guidelines and undertake a qualified assessment considering:

- identified health effects of each pollutant;
- likely emission sources; and
- potentially exposed populations.

<sup>21</sup> NB: Health incidence data are specific to the census area units under assessment (refer Figure 1)

## 4. QUANTIFIED CHRONIC EXPOSURE: PM<sub>10</sub>, PM<sub>2.5</sub> & NO<sub>2</sub> IN MOUNT MAUNGANGUI

---

### 4.1 PM<sub>10</sub>

#### Otūmoetai

The monitoring station is located at Otūmoetai Primary School which is located in the Otūmoetai South census area unit. The area is residential, and concentrations measured at this location should be representative of the adjacent census area units. This value was therefore adopted as representative of Otūmoetai and surrounding residential suburbs (Otūmoetai South, Otūmoetai North, Bellevue & Matua census area units) for the base year 2019.

Otūmoetai has no annual average PM<sub>10</sub> data for the year 2019 due to instrument failure. However, annual PM<sub>10</sub> concentrations measured at this location have been stable, averaging 10 µg/m<sup>3</sup> for the eight years ending 2021 (max 11, min 9, std dev 0.5).

An annual average concentration of 10 µg/m<sup>3</sup> was assumed for PM<sub>10</sub> for the year 2019 for all Otūmoetai census area units.

#### Mount Maunganui

**Figure 8** shows the BOPRC PM<sub>10</sub> monitoring site locations in the Mount Maunganui Airshed.

Ambient air quality monitoring is undertaken in three of the five census area units under assessment, notably Mount Maunganui North (Rata Street), Tauranga City Marinas (Tauranga Bridge Marina) and Sulphur Point. We used the PM<sub>10</sub> mean concentration from 2019 measured at these monitoring site locations as the key metric for exposure in these census area units.

For the two remaining census areas units, Omanu and Arataki, it is important to consider spatial variation. The Totara Street monitoring site is in the middle of the industrial area and measurements at this location may not be representative of long-term concentrations in other parts of Omanu, or Arataki.

**Figure 8** includes a wind rose for the year 2019, overlaid on the monitoring locations in the Mount Maunganui Airshed. The wind rose shows the predominant wind directions are (from the) south-west quadrant. This means that, over the year, emissions from the industrial and port activities will be directed towards to the north-east more than other locations.


This directional trend is evident when comparing annual PM<sub>10</sub> concentrations measured on the south-west boundary of the Mount Maunganui Airshed (Sulphur Point, Bridge Marina and Whareroa Marae) with those measured inside the airshed and to the north (Rata Street, Rail Yard South and Totara Street refer **Table 5**).<sup>22</sup> Annual concentrations are ~60% higher, on

---

<sup>22</sup> Long-term PM<sub>10</sub> measured at De Havilland Way is dominated by adjacent industrial sources and excluded from this consideration.

average, inside and to the north of the Mount Maunganui Airshed compared with monitoring locations on the predominantly upwind, south-west boundary.

**FIGURE 8: 2019 PM<sub>10</sub> monitoring locations and annual PM<sub>10</sub> concentrations measured in the Mount Maunganui Airshed** [Source: BOPRC. Insert 2019 wind rose from Whareroa Marae (refer Fig A10)]



**TABLE 5: Comparison of mean 2019 PM<sub>10</sub> concentrations at Mount Maunganui monitoring locations**

South-West Boundary Monitoring Site	2019 PM <sub>10</sub> (µg/m <sup>3</sup> )	Northern / Inside Airshed Monitoring Site	2019 PM <sub>10</sub> (µg/m <sup>3</sup> )
Sulphur Point	14	Rata Street	20
Tauranga Bridge Marina	16	Rail Yard South	31
Whareroa Marae	17	Totara Street	25
		De Havilland Way	20
<b>Average:</b>	<b>15</b>	<b>Average:</b>	<b>24</b>
<b>Difference: 8 (55%)</b>			

It is also notable, however, that mean 2019 PM<sub>10</sub> concentrations measured on the south-west boundary of the Mount Maunganui Airshed (Sulphur Point, Bridge Marina and Whareroa Marae average 15 µg/m<sup>3</sup>) are elevated in comparison with the residential suburb of Otūmoetai (10 µg/m<sup>3</sup>).

The Rata Street monitoring location was selected as indicative of long-term PM<sub>10</sub> levels in Omanu & Arataki on the basis that:

- Mean 2019 concentrations measured at Totara Street and Rail Yard South were impacted by adjacent industrial & port activities and are too high to be representative of Omanu & Arataki;
- The Rata Street monitoring site is in a residential location that is often downwind of emissions from the Port activities;
- The De Havilland Way monitoring site is adjacent to residential activity with the same mean 2019 PM<sub>10</sub> value (20 µg/m<sup>3</sup>) as Rata Street;
- There will necessarily be some attenuation with distance, although how much is uncertain; and
- Whilst the Rata Street location may underestimate long-term PM<sub>10</sub> levels in Omanu & Arataki, this serves to ensure that the estimate of associated health effects is not an overestimate.

**Table 6** presents mean PM<sub>10</sub> concentrations assigned to each census area unit for assessment purposes.

**TABLE 6: Mean 2019 PM<sub>10</sub> concentrations assigned to each census area unit**

Census Area Unit	2019 mean PM <sub>10</sub> (µg/m <sup>3</sup> )	Reference
Otūmoetai South	10	Otūmoetai monitoring station
Otūmoetai North	10	Otūmoetai monitoring station
Bellevue	10	Otūmoetai monitoring station
Matua	10	Otūmoetai monitoring station
Mount Maunganui North	20	Rata Street monitoring station
Omanu	20	Rata Street monitoring station
Tauranga City-Marinas	16	Tauranga Bridge Marina monitoring station
Arataki	20	Rata Street monitoring station
Sulphur Point	14	Sulphur Point monitoring station

## 4.2 PM<sub>2.5</sub>

### Otūmoetai

There is no PM<sub>2.5</sub> monitoring station in Otūmoetai. PM<sub>2.5</sub> is a component of PM<sub>10</sub> and the two size fractions will correlate with each other. Kuschel *et al.*,(2022) assigned a ratio of 0.51 for



PM<sub>2.5</sub>:PM<sub>10</sub> in Otūmoetai South based on an empirical relationship with biomass. We similarly used this ratio to estimate mean 2019 PM<sub>2.5</sub> concentrations from available PM<sub>10</sub> data.

A PM<sub>2.5</sub> annual average concentration of 5.3 µg/m<sup>3</sup> was estimated for the year 2019 for all Otūmoetai census area units.

### Mount Maunganui

There is only one PM<sub>2.5</sub> monitoring station in the Mount Maunganui area, located at Totara Street in the Mount Maunganui Airshed.

PM<sub>2.5</sub> is a component of PM<sub>10</sub> and the two size fractions will correlate with each other (noting that this correlation differs between industrial & residential settings). The ratio of (annual) PM<sub>2.5</sub> to PM<sub>10</sub> measured at Totara Street in 2019 was 0.32. This is similar to PM<sub>2.5</sub> to PM<sub>10</sub> ratios measured in other industrial settings (Sridhar *et al.*, 2022).

The mean 2019 Totara Street PM<sub>2.5</sub>:PM<sub>10</sub> ratio was used to estimate mean 2019 PM<sub>2.5</sub> concentrations for all Mount Maunganui census area units based on their previously estimated PM<sub>10</sub> data. This may underestimate PM<sub>2.5</sub> in these locations, as PM<sub>2.5</sub> in residential areas is typically higher (fraction of PM<sub>10</sub>). However, it serves to ensure the estimate does not overestimate concentrations of PM<sub>2.5</sub> and associated health effects.

**Table 7** presents mean 2019 PM<sub>2.5</sub> concentrations assigned to each census area unit for assessment purposes.

**TABLE 7: Mean 2019 PM<sub>2.5</sub> concentrations assigned to each census area unit**

Census Area Unit	2019 mean PM <sub>2.5</sub> (µg/m <sup>3</sup> )	Reference
Otūmoetai South	5.3	2019 PM <sub>10</sub> Otūmoetai, ratio 0.51 (Sridhar et al., 2022)
Otūmoetai North	5.3	2019 PM <sub>10</sub> Otūmoetai, ratio 0.51 (Sridhar et al., 2022)
Bellevue	5.3	2019 PM <sub>10</sub> Otūmoetai, ratio 0.51 (Sridhar et al., 2022)
Matua	5.3	2019 PM <sub>10</sub> Otūmoetai, ratio 0.51 (Sridhar et al., 2022)
Mount Maunganui North	6.4	2019 PM <sub>10</sub> Rata St, ratio 0.32 (Totara St, 2019)
Omanu	6.4	2019 PM <sub>10</sub> Rata St, ratio 0.32 (Totara St, 2019)
Tauranga City-Marinas	5.1	2019 PM <sub>10</sub> Bridge Marina, ratio 0.32 (Totara St, 2019)
Arataki	6.4	2019 PM <sub>10</sub> Rata St, ratio 0.32 (Totara St, 2019)
Sulphur Point	4.5	2019 PM <sub>10</sub> Sulphur Point, ratio 0.32 (Totara St, 2019)

### 4.3 NO<sub>2</sub>

Concentrations assigned to each census area unit in the HAPINZ 3 model for a base year 2016 were updated to 2019 by applying a scalar of 0.9354 to all locations as recommended in Kuschel *et al.*, 2022.

NB: This will not assess peak NO<sub>2</sub> concentrations likely to be experienced by residents living (and working) close to roads and motorways. Roadside monitoring by Waka Kotahi in the

Mount Maunganui Airshed<sup>23</sup> indicates significantly higher annual average levels in 2018-2019 (34 µg/m<sup>3</sup>) to that assigned for the Omanu census area unit (6 µg/m<sup>3</sup>).

**Table 8** presents mean 2019 NO<sub>2</sub> concentrations assigned to each census area unit for assessment purposes.

**TABLE 8: Mean 2019 NO<sub>2</sub> concentrations assigned to each census area unit**

Census Area Unit	2019 mean NO <sub>2</sub> (µg/m <sup>3</sup> )
Otūmoetai South	6.0
Otūmoetai North	7.9
Bellevue	6.0
Matua	5.1
Mount Maunganui North	7.8
Omanu	6.2
Tauranga City-Marinas*	12
Arataki	8.1
Sulphur Point	12

\* Tauranga City-Marinas not assigned NO<sub>2</sub> value in HAPINZ 3 – value from Sulphur Point assumed

<sup>23</sup> HAM008 Maunganui Rd / Golf Rd12

## 5. QUANTIFIED HEALTH IMPACTS: PM<sub>10</sub>, PM<sub>2.5</sub> & NO<sub>2</sub> IN MOUNT MAUNGANGUI

### 5.1 PM<sub>2.5</sub> AND NO<sub>2</sub>

**Table 9** presents estimated premature mortality and morbidity associated with long-term exposure to PM<sub>2.5</sub> and NO<sub>2</sub> in Mount Maunganui and Otūmoetai for a base year of 2019.

**TABLE 9: Estimated premature mortality and morbidity associated with annual PM<sub>2.5</sub> and NO<sub>2</sub>**

Health Effect	Mount Maunganui (5xCAU)	Otūmoetai (4xCAU)	Difference (Δ)
Population (2019)	16,975	17,835	860
<b>Cases due to PM<sub>2.5</sub></b>			
Premature death (adults 30 years+)	9	7	2
Cardiovascular hospitalisations (all ages)	16	13	3
Respiratory hospitalisations (all ages)	11	9	2
Restricted activity days (all ages)	9,764	8,507	1,256
Cost (2019 \$NZD)	\$43.2 M	\$33.6 M	\$9.6 M
<b>Cases due to NO<sub>2</sub></b>			
Premature death (adults 30 years+)	10	7	3
Cardiovascular hospitalisations (all ages)	8	7	1
Respiratory hospitalisations (all ages)	23	19	4
Asthma prevalence (0-18 years)	40	38	2
Cost (2019 \$NZD)	\$46.4 M	\$34.0 M	\$12.4 M
<b>Cases due to PM<sub>2.5</sub> and NO<sub>2</sub></b>			
Premature death (adults 30 years+)	19	14	5
95% Confidence intervals	(14, 25)	(10, 19)	(3, 6)
Cardiovascular hospitalisations (all ages)	24	20	4
Respiratory hospitalisations (all ages)	34	28	6
Restricted activity days (all ages) (PM <sub>2.5</sub> )	9,764	8,507	1,256
Asthma prevalence (0-18 years) (NO <sub>2</sub> )	40	38	2
Cost (2019 \$NZD)	\$90 M	\$68 M	\$22 M

The population in the areas assessed are very similar (~17,000), with Otūmoetai having slightly more (860) people. Thus, the difference between estimates of health impacts for each area is assumed to be attributable to the difference in exposure to air pollution.



The modelling estimates that the Mount Maunganui area has around five premature deaths each year associated with increased exposure to long-term concentrations of PM<sub>2.5</sub> and NO<sub>2</sub> when compared with Otūmoetai. For context, the total mortality from all non-external causes in Mount Maunganui for the year 2019 was 145 so this estimate represents around 3% of deaths in that year.

## 5.2 PM<sub>10</sub>

**Table 10** presents estimated premature mortality associated with long-term exposure to PM<sub>10</sub> in Mount Maunganui and Otūmoetai for a base year of 2019. The population in the areas assessed are very similar (~17,000), with Otūmoetai having slightly more (860) people. Thus, the difference between estimates of health impacts for each area can be assumed to be primarily attributable to the difference in exposure to air pollution.<sup>24</sup>

The modelling estimates that the Mount Maunganui area has 13 premature deaths (95% confidence interval 11 to 15) each year associated with increased exposure to long-term concentrations of PM<sub>10</sub> when compared with Otūmoetai. For context, the total mortality from all non-external causes<sup>25</sup> in Mount Maunganui for the year 2019 was 145 so this estimate represents around 9% of deaths in that year.

**TABLE 10: Estimated premature mortality associated with annual PM<sub>10</sub>**

Health Effect	Mount Maunganui (5xCAU)	Otūmoetai (4xCAU)	Difference (Δ)
Population (2019)	16,975	17,835	860
	<b>Cases due to PM<sub>10</sub></b>		
Premature death (adults 30 yrs+)	26	13	13
95% Confidence intervals	(22, 31)	(11, 16)	(11,15)
Cost (2019 \$NZD)	\$119 M	\$61 M	\$59 M

## 5.3 MEETING THE WHO AQG 2021 GUIDELINES

Toi Te Ora Public Health requested assessment of the health impacts avoided if all areas met the WHO 2021 global air quality guidelines. The hypothetical scenario of all census area units were modelled, including Otūmoetai, meeting an annual PM<sub>2.5</sub> guideline of 5 µg/m<sup>3</sup> and an annual NO<sub>2</sub> guideline of 10 µg/m<sup>3</sup>. **Table 11** presents the health impacts avoided if the WHO guidelines were met in all areas (i.e., both Otūmoetai and Mount Maunganui).

All areas except Tauranga Bridge Marina and Sulphur Point, both of which have very low usually resident populations, already meet the WHO 2021 global air quality guideline for NO<sub>2</sub>. Consequently, the improvements outlined in **Table 11** would be entirely due to (hypothetical) reductions in annual PM<sub>2.5</sub>.

<sup>24</sup> Assuming base health incidence rates in each area are similar (mean 2019 all-cause mortality was 8% higher in Mount Maunganui)

<sup>25</sup> i.e., deaths excluding accidents and violence



**TABLE 11: Estimated avoided health impacts and social costs if WHO guidelines met in Otūmoetai and Mount Maunganui versus base case**

Health Effect	2019 Air Quality	Meet WHO AQG	Δ
Premature death (adults 30 years +)	34	31	-2
Cardiovascular hospitalisations (all ages)	43	39	-4
Respiratory hospitalisations (all ages)	63	60	-3
Restricted activity days (all ages)	18,271	15,663	-2,608
Asthma Prevalence (0-18 years)	78	78	0
Cost (2019 \$NZD)	\$157 M	\$146 M	-\$11 M

#### 5.4 SENSITIVITY TESTING

The HAPINZ 3 Health Effects Model includes 95% confidence intervals for the (New Zealand specific) exposure response functions and value of statistical life as well as low and high-end social cost estimates. Using these ranges, we tested the sensitivity of the model estimates for PM<sub>2.5</sub> and NO<sub>2</sub>.

**Table 12** summarises the results of the sensitivity testing in terms of social costs for the modelling of effects associated with PM<sub>2.5</sub> and NO<sub>2</sub>.

NB: The base case under consideration is the difference (Δ) in estimates between Mount Maunganui and Otūmoetai as this the fraction that can be attributed to anthropogenic emissions from industrial and port activities (PM<sub>2.5</sub> and NO<sub>2</sub>), and transport (NO<sub>2</sub>).

Using the high estimates for the social cost values would have the greatest impact (+34%) on total anthropogenic social costs.

**TABLE 12: Sensitivity of the HAPINZ 3 social cost estimates for PM<sub>2.5</sub> and NO<sub>2</sub> to different parameters versus the base case (in 2019 NZD)**

Scenario	Social Cost (2019 \$million)			
	Δ Mortality	Δ Morbidity	Δ Total	vs base case
<b>Base case</b>				
Default values	21.5	0.4	22	-
<b>Exposure response function values</b>				
Lower bound 95% CI	15.5	0.3	16	-28%
Upper bound 95% CI	27.1	0.6	28	+26%
<b>Social cost values</b>				
Lower estimate	19.3	0.1	19	-12%
High estimate	24.9	4.5	29	+34%

#### 5.4.1 Effect of exposure response function values

If the **lower bounds of the confidence intervals** for the exposure-response functions were used, then the impacts would be 28% less than those in the base case.

If the **higher bounds of the confidence intervals** for the exposure-response functions were used, then the impacts would be 26% higher than those predicted using the base case.

**Overall effect:** Within these two bounds, the social cost of anthropogenic air pollution health impacts are between \$16 million and \$28 million (relative to the base case of \$22 million) in Mount Maunganui as compared with Otūmoetai.

Note: The number of premature deaths for adults due to anthropogenic air pollution (PM<sub>2.5</sub> and NO<sub>2</sub>) in Mount Maunganui as compared with Otūmoetai would be between 3 and 6 (relative to the base case of 5 deaths).

#### 5.4.2 Effect of social cost values

If the **low estimates** for the social cost values were used, then the impacts would be 12% less than those in the base case.

If the **high estimates** for the social cost values were used, then the impacts would be 34% more than those in the base case.

**Overall effect:** Within these two bounds, anthropogenic air pollution health impacts in Mount Maunganui as compared with Otūmoetai are between \$19 million and \$29 million (relative to the base case of \$22 million).

Note: The increase in the high-cost estimate scenario is largely due to factoring in loss of life quality from prolonged illness and suffering which is not included in the base case. Many overseas jurisdictions use a specific environmental value of statistical life which is considerably higher than a transport (road safety) risk value to adequately account for these loss-of-life quality costs.

### 5.5 UNCERTAINTY & CONFIDENCE ASSESSMENT

The key sources of uncertainty in this assessment of air pollution health impacts are described as follows.

#### Air pollutants exist as a complex mixture

There is a considerable body of evidence from epidemiological studies confirming the adverse health effects associated with exposure to air pollution. Notably, WHO determined all PM-outcome associations were deemed causal, or likely to be causal (WHO, 2021). However, the adverse effects attributed to nitrogen dioxide may actually be attributable to other pollutants in the mixture.

#### Baseline disease burden

Data on the number of deaths and cases of disease can be uncertain, particularly if data from a number of sources are combined or if projections of future cases are made. In this study, health incidence statistics in the HAPINZ 3.0 model for a base year 2016 were updated with Ministry of Health statistics for a base year 2019 (Metcalf & Kuschel, pending).

We have a high degree of confidence in the data.

## Pollution exposure level

This is a potentially significant limitation of the modelling. The approach in this study was to estimate long-term exposure based on:

- Available measured pollutant concentrations at representative monitoring locations for PM<sub>10</sub>;
- An assigned ratio (for PM<sub>2.5</sub>:PM<sub>10</sub>) based on an empirical relationship with biomass (Kuschel *et al.*, 2022) to establish PM<sub>2.5</sub> in Otūmoetai;
- A measured ratio (for PM<sub>2.5</sub>:PM<sub>10</sub>) to establish PM<sub>2.5</sub> in Mount Maunganui;
- Spatial estimates for NO<sub>2</sub> in the national model updated for base year 2019 by applying a scalar as recommended in Kuschel *et al.*, 2022.

The general approach in estimating long-term levels of key pollutants was one of caution to avoid overstating potential impacts. This means the estimates are not conservative (i.e., likely underestimates).

The base year of 2019 pre-dates potential impacts of the pandemic on ambient air quality pollutants in the Mount Maunganui and Otūmoetai areas.

We estimate the exposure assessment is within +/-25%.

## The exposure-response function

Exposure response functions are derived from epidemiological studies, in which assumptions made in the analysis inevitably introduce some uncertainty into the results. The HAPINZ 3.0 model utilises New Zealand specific exposure-response functions for NO<sub>2</sub> and PM<sub>2.5</sub>. This means that the uncertainty in the exposure assessment is captured (to some extent) in the uncertainty of the exposure-response functions (and represented in the quoted 95% confidence intervals).

## The counterfactual level of air pollution

The counterfactual level of air pollution is the baseline or reference exposure against which the health impacts of air pollution are calculated (e.g., having no air pollution). This is not a source of uncertainty in itself.

The modelling was comparative, i.e., only the difference between long-term exposure in Mount Maunganui and Otūmoetai was assessed. As such the modelling estimates are not sensitive to the counterfactual.

## Confidence Assessment

Hales *et al.*, assessed the risk of bias in accordance with WHO, 2020 and concluded it was low for all factors except potential confounding due to the inability to control for BMI due to a lack data in NZ health incidence statistics. This increased the overall risk of bias to low-to-moderate.

The HAPINZ 3.0 model has been internationally peer reviewed and warrants a high degree of confidence for the base case 2016 (Kuschel *et al.*, 2022).

The HAPINZ 3.0 model has been updated with 2018-2019 air quality, population and health incidence data which increases its representativeness. Significantly higher uncertainties are

associated with the assumptions used to estimate pollution exposure. Cumulatively, the overall uncertainty associated with the estimates is around +/-30%.

Overall, we have a moderate degree of confidence in the model estimates.

## **5.6 MODEL LIMITATIONS**

It should be noted that the quantified assessment relies on a published model (HAPINZ 3.0) to assess the specified health outcomes for chronic exposure to PM<sub>2.5</sub>, NO<sub>2</sub> or PM<sub>10</sub> only. Other potential adverse health outcomes associated with exposure to chronic levels of other pollutants that may be present in Mount Maunganui are not, and cannot be, considered. Such effects could include, for example, reproductive fertility, developmental toxicity and cancer risk (e.g., known benzene specific cancers).

## 6. QUALITATIVE ASSESSMENT: SO<sub>2</sub>, H<sub>2</sub>S, BENZENE & ODOUR IN MOUNT MAUNGANGUI

---

The above quantified assessment of PM<sub>10</sub>, PM<sub>2.5</sub> and NO<sub>2</sub> using internationally peer reviewed, quantified risk ratios has inevitably focussed attention on PM and NO<sub>2</sub>. This provides a good indication of overall health impacts due to exposure to air pollution from industry, motor vehicles and domestic fires in Mount Maunganui as compared with Otūmoetai.

However, residents of Mount Maunganui are also exposed to elevated levels of other contaminants from location-specific industries. For example, ambient concentrations of SO<sub>2</sub> are known to be elevated in some locations due to emissions from shipping and manufacturing. Impacts from these pollutants are not addressed in the above quantified assessment. This section addresses other pollutants known or suspected to be elevated in and around Mount Maunganui.

It is important to note that total ship visits to the Port of Tauranga dropped by 20% in 2020 and 2021, compared with 2018 and 2019, due to the global COVID-19 pandemic.<sup>26</sup> Overall industrial activity and associated vehicle movements in the Mount Maunganui Airshed were also reduced during intermittent lockdown periods during this time. The scale and duration of reduced activity, and any associated reduction in emissions, is not known.

### 6.1 SULPHUR DIOXIDE

#### 6.1.1 Air quality exposure to SO<sub>2</sub>

**Figure 8** shows the BOPRC SO<sub>2</sub> monitoring site locations in the Mount Maunganui Airshed.

**Appendix A** (Tables A1-A9) presents summary data for 10-minute, hourly and daily SO<sub>2</sub> monitoring in the Mount Maunganui Airshed for the years 2019 – 2021 respectively for comparison with health-based standards and guidelines.

Also in **Appendix A** are:

- Time series ambient air quality data are presented graphically in for 10-minute, hourly and daily concentrations for the years 2019 – 2021 respectively (Figures A1 – A9); and
- Frequency of wind direction (from) and wind speeds (wind roses) measured at Whareroa Marae for the years 2019 – 2021 respectively (Figures A10 – A13). These show that winds are predominantly from the southwest quadrant (nearly ~30%), and to a lesser extent (just under ~10%) from the northeast.

#### Discussion

Monitoring at Whareroa Marae commenced in September 2015 following repeated public complaints to BOPRC regarding *inter alia* poor air quality and wide-ranging respiratory issues, watering eyes and sore throats.

---

<sup>26</sup> Difference calculated from average total ship visits in 2018 and 2019 compared with the average total ship visits in 2020 and 2021.

Our review of this monitoring reveals multiple breaches of the national environmental standard for SO<sub>2</sub> (upper limit 570 µg/m<sup>3</sup>, zero permitted exceedances). These breaches were attributed to the fertiliser works, which is located adjacent to Whareroa Marae, by BOPRC who undertook regulatory action in 2016. The plant significantly reduced its emissions and ambient levels of SO<sub>2</sub> measured at Whareroa Marae have not breached national environmental standards since 2016.

In late 2018, continuous, ambient air quality monitoring data commenced at six other locations in **Figure 8**. As evident from the data summary and time series graphs, the data show intermittent, elevated short-term levels of sulphur dioxide continued to occur at all locations throughout 2019.

A dramatic reduction in ambient levels of SO<sub>2</sub> appears to coincide with ships (registered overseas) implementing Annex VI of MARPOL on 1 January 2020.<sup>27</sup> Annex VI of MARPOL requires ships to burn low sulphur fuel or implement abatement technology to mitigate emissions of SO<sub>2</sub>.

This reduction was most evident at monitoring sites located closest to (i.e., within 500 metres), and predominantly downwind of, the berths; namely Rata Street and Sulphur Point, Rail Yard South and Totara Street (refer graphs of 10-minute SO<sub>2</sub>, Figure A1 and A2, 1-hour SO<sub>2</sub>, Figure A4 and A5 and daily SO<sub>2</sub>, Figure A7 and A8). Since 1 January 2020 there have been no exceedances of either the 10-minute (2006) or daily (2021) WHO guidelines for SO<sub>2</sub> at these locations, whereas prior to 1 January 2020, exceedances of these guidelines occurred regularly.

Whareroa Marae and Tauranga Bridge Marina monitoring sites, which are residential/mixed use locations, continue to record occasional exceedances of WHO 10-minute and daily guidelines. We have carefully reviewed this data for monthly and annual reporting to Toi Te Ora and noted temporal coincidence of elevated concentrations with wind directions from the adjacent fertiliser works towards these monitoring sites.<sup>28</sup> The occasional exceedances of the WHO 10-minute and daily SO<sub>2</sub> guidelines at these two locations appear to be influenced primarily by SO<sub>2</sub> emissions from the fertiliser works.<sup>29</sup>

### 6.1.2 Assessment: SO<sub>2</sub>

Ambient levels of SO<sub>2</sub> above the WHO 10-minute guideline may have resulted in some short-term, transient effects at locations such as Whareroa Marae and Tauranga Bridge Marina including:

- Bronchoconstriction (particularly asthmatics when exercising); and/or
- Nose or throat irritation

---

<sup>27</sup> New Zealand acceded to Annex VI in late 2021 so this reduction likely reflects the majority of ships visiting the Port of Tauranga being registered in countries that have already ratified Annex VI.

<sup>28</sup> For example, Memo from L. Wickham (EIL) to S. Halligan (MoH) dated 7 April 2022 Re: *Three-year review (2019-2021) of air quality monitoring at Mount Maunganui*.

Memo from L. Wickham (EIL) to C. Lochore (Toi Te Ora) dated 12 March 2021 Re: *Two-year review (2019/2020) of air quality monitoring at Mount Maunganui*.

Memo from L. Wickham (EIL) to S. Layne & J. Miller (Toi Te Ora) dated 7 February 2020. Re: *Annual review: 2019 air quality monitoring at Mount Maunganui*.

<sup>29</sup> NB: Exceedance of a WHO (global) air quality guideline has no regulatory status in New Zealand. This is different to a 'breach' of a national environmental standard for SO<sub>2</sub> (which previously led to regulatory action by BOPRC as noted above).

Those most likely to experience effect would be asthmatics, babies and infants, children and elderly people. However, being transient, any effects are likely to have been short-lived (i.e., over within minutes or hours of ambient sulphur dioxide levels returning to normal).

It is less clear what the effects of continued exceedances of the WHO daily guideline would be. Recent epidemiological evidence supports daily increases in SO<sub>2</sub> being causal for increased risk of all-cause mortality and respiratory mortality (Orellano *et al.*,2021) with a high certainty of evidence (WHO, 2021). This would not be reflected in the quantified mortality estimates generated separately for PM<sub>10</sub> and PM<sub>2.5</sub> and NO<sub>2</sub>. Recent epidemiological evidence further correlated daily increases in SO<sub>2</sub> with increased risk of asthma-associated emergency room visits and hospital admissions (Zheng *et al.*,2021) and this relationship is causal (WHO, 2021).

The daily levels of SO<sub>2</sub> in the referenced studies above are not dissimilar to daily levels of SO<sub>2</sub> measured at Whareroa Marae and Tauranga Bridge Marina (Appendix A). No threshold for effects has been identified (WHO, 2021).

The evidence suggests that people living at Whareroa Marae and the Tauranga Bridge Marina (on boats) may have been, and continue to be, adversely affected by SO<sub>2</sub> emissions. The focus in this assessment has been on chronic exposure and we have not quantified these effects.


## 6.2 HYDROGEN SULPHIDE

### 6.2.1 Air quality exposure to H<sub>2</sub>S

BOPRC monitored H<sub>2</sub>S at Whareroa Marae between September 2015 and October 2020, after which time monitoring ceased. **Figure 9** presents a time series of hourly H<sub>2</sub>S data during this period and **Table 12** presents summary statistics. **Table 12** shows that the New Zealand guideline for H<sub>2</sub>S was regularly exceeded at Whareroa Marae.



**FIGURE 9: Hourly H<sub>2</sub>S concentrations measured at Whareroa Marae; 26 Sep 2015 – 14 Oct 2020** [Source: BOPRC]



**TABLE 13: Summary hourly H<sub>2</sub>S concentrations at Whareroa Marae 2016 – 2020**

Year	Maximum	99 <sup>th</sup> Percentile	Standard Deviation	No. exceedances NZAAQG (7 µg/m <sup>3</sup> )
	(µg/m <sup>3</sup> , 1-hour average)			
2016	14	3	1	81
2017	36	4	2	100
2018	14	3	1	22
2019	17	4	1	35
2020*	19	4	1	54

\*Partial year (to 14 Oct only)

### 6.2.2 Assessment: Hydrogen Sulphide

The monitoring record at Whareroa Marae shows many exceedances of the New Zealand H<sub>2</sub>S guideline that was set to protect against offensive odours. A detailed source apportionment by BOPRC supports several industrial sources to the north of Whareroa Marae as responsible, with odours from natural sources possibly playing a part during summer (BOPRC 2020).

There is an established history of odour complaint from the marae to BOPRC (BOPRC, 2020). We note that Te Rūnanga o Ngai Te Rangi Iwi Trust have provided evidence to the Environment Court stating tangata whenua have experienced adverse cultural impacts due to industrial odours.<sup>30</sup>

We conclude that industrial emissions of H<sub>2</sub>S have regularly caused offensive odours at Whareroa Marae. This would be reducing the quality of life and impacting adversely on the wellbeing of residents and visitors, including manuhiri (guests).

### 6.3 BENZENE

Benzene is discharged to air from motor vehicles as well as industrial sources in the Mount Maunganui area. Of note are the presence of multiple bulk fuel storage facilities at the port that will have fugitive emissions of benzene. Emissions of benzene have not been quantified or assessed, however, given the presence of an oil re-refinery and the number of large fuel tanks present (Refer **Figure 10**), ambient concentrations of benzene may be significant.

Background levels of benzene in areas with no vehicle or industry are typically very low (< 1 ppb, WHO 2000).

<sup>30</sup> Statement of Evidence of Mr Reon Tuanau dated 14 August 2020. At [30]. ENV-2019-AKL-000065.

**FIGURE 10: Benzene sources Mount Maunganui (traffic, fuel tanks, oil re-refinery) in relation to Whareroa Marae & Tauranga Bridge Marina**



### 6.3.1 Air quality exposure to benzene

#### Acute

There has been limited ambient monitoring of benzene in the Mount Maunganui Airshed. Two industry reports include monitoring of ambient benzene (i.e., beyond the site boundary) in industrial locations of the Mount Maunganui Airshed (Ecocific 2020, 2021). These consistently measured ambient concentrations of benzene above the OEHHA REL for benzene as shown in **Table 14**. The maximum concentration was 130 ppb as a 1-hour average, measured in a location that was within 150 metres of a tank farm. However, some of the data may not be robust (lower detection limit reported as 70 ppb). Monitoring in an industrial location in Christchurch measured similar 1-hour average concentrations of benzene (8 – 93 ppb, n = 8).<sup>31</sup>

<sup>31</sup> Specifically downwind of a wastewater treatment plant under upset conditions. Source: Christchurch City Council, unpublished data

**TABLE 14: Hourly benzene concentrations measured at Mount Maunganui, 2020-2021**

Ecocific 2020, 2021	Benzene (ppb, 1-hour average)
OEHHA REL	8
Maximum	130
Average	47
Minimum	10
n*	12

\*Number of hourly measurements

## Chronic

We are not aware of any long-term monitoring of benzene in Mount Maunganui. This precludes comparison with the New Zealand long-term ambient air quality guideline or quantification of risk due to carcinogenicity.

Long-term ambient monitoring in Auckland suggests people living close to busy roads (~13,000 vehicles per day) and petrol stations may be exposed to annual concentrations of benzene of around 0.6 - 1 ppb respectively (Auckland Council, 2014). A 2012 literature review prepared for Auckland Council (Emission Impossible Ltd, 2012) concluded annual concentrations of benzene may be elevated within 150 metres of busy roads, but this is of uncertain relevance for Mount Maunganui in 2022.

### 6.3.2 Assessment: acute benzene

There are several potentially significant sources of benzene in the Mount Maunganui Airshed, however, these have not been quantified or comprehensively assessed to date. Short-term ambient monitoring for benzene in the Mount Maunganui Airshed has recorded 1-hour average concentrations above the OEHHA reference exposure level (8 ppb) set to protect against developmental effects. These measurements were undertaken in industrial locations however, they were ambient measurements, i.e., beyond the boundary of the industrial site that was undertaking the measurement. The monitoring suggests that workers in these locations may have elevated, acute exposure to benzene. However, the monitoring data are limited (12 hours sampling over two years only) and in the absence of an emissions inventory or more detailed assessment, no firm conclusions can be drawn.

It is further unclear if, and how, these elevated acute measurements relate to residential exposure in other parts of the Mount Maunganui Airshed (e.g., Whareroa Marae, Rata Street). Whareroa Marae is within 200 metres of a bulk fuel storage facility and within 300 metres of an oil re-refining facility. Residents and visitors (including manuhiri) to Whareroa Marae and kohanga reo may be exposed to elevated concentrations of benzene over acute and/or chronic time frames. Benzene exposure may disproportionately impact infants and children (OEHHA, 2014).

There are no long-term monitoring data for benzene in the Mount Maunganui Airshed. This is a significant data gap. No conclusions can be drawn about either worker or residential chronic exposure to benzene.

## 6.4 ODOUR

Odour is an established issue for Mount Maunganui. In the year ending 1 September 2021, BORC received 512 odour complaints to their pollution hotline for the Mount industrial area (BOPRC, undated). The majority of these complaints were about pet food, bitumen, and 'rotten egg' (H<sub>2</sub>S).

The large number of industrial sources and the significance of the odour issues in the Mount Maunganui Airshed (more than 500 complaints a year) are such that assessment of any, single, "indicator" compound (e.g., hydrogen sulphide) would be insufficient to understand the extent of potential public health impacts from odour. This is because odour is made up multiple contaminants.

It is apparent that offensive and objectionable odours are reducing the quality of life and adversely impacting on the wellbeing of residents in and around the Mount Maunganui Airshed.

A nascent body of epidemiology suggests that industrial odours are associated with adverse health impacts in surrounding communities (e.g., Guadalupe-Fernandez *et al.*, 2021). This suggests that, in addition to negative impacts on wellbeing, odorous emissions may also be adversely impacting residents' health.

## 7. CONCLUSIONS

---

Our review of ambient air quality monitoring data shows that concentrations of some air contaminants are elevated in some locations relative to health and wellbeing-based ambient criteria.

### Quantitative Assessment

The modelling estimates that, compared with Otūmoetai, in Mount Maunganui there were:

- Around five premature deaths each year associated with increased exposure to long-term concentrations of **PM<sub>2.5</sub>** and **NO<sub>2</sub>**. For context, the total mortality from all non-external causes<sup>32</sup> in Mount Maunganui for the year 2019 was 145 so this estimate represents around 3% of deaths in that year.
- An additional four cardiovascular and six respiratory hospitalisations associated with increased long-term exposure to **PM<sub>2.5</sub>** and **NO<sub>2</sub>**.
- An additional 1,256 restricted activity days associated with increased long-term exposure to **PM<sub>2.5</sub>**.
- Two additional cases of asthma in under 18-year-olds associated with increased long-term exposure to **NO<sub>2</sub>**.

Sensitivity testing estimates that:

- The number of premature deaths for adults associated with long-term exposure to **PM<sub>2.5</sub>** and **NO<sub>2</sub>** in Mount Maunganui as compared with Otūmoetai have a 95% confidence interval of 3 and 6 (relative to the base case of 5 deaths).
- The low and high social cost estimates associated with long-term exposure to **PM<sub>2.5</sub>** and **NO<sub>2</sub>** in Mount Maunganui as compared with Otūmoetai are between \$19 million and \$29 million (relative to the base case of \$22 million, all in NZ\$2019).

Single-pollutant modelling estimated that the Mount Maunganui area has 13 premature deaths each year (95% confidence interval 11 to 15) associated with increased exposure to long-term concentrations of **PM<sub>10</sub>** when compared with Otūmoetai. This estimate represents around 9% of total mortality from all non-external causes in that year, which is higher than the estimates associated with long-term exposure to **PM<sub>2.5</sub>** and **NO<sub>2</sub>**. It should be noted that the **PM<sub>10</sub>** modelling is not additive to the estimate of effects associated with **PM<sub>2.5</sub>** and **NO<sub>2</sub>**, rather it is a different estimate.

At Toi Te Ora Public Health's request, we modelled the hypothetical scenario of all areas, including Otūmoetai, meeting the annual WHO 2021 global air quality guidelines for **PM<sub>2.5</sub>** (5 µg/m<sup>3</sup>) and **NO<sub>2</sub>** (10 µg/m<sup>3</sup>). This would have a modest impact in averting 2 premature deaths per year with an associated averted social cost of \$11 million (NZ\$2019) due to hypothetical reductions in annual **PM<sub>2.5</sub>** (as our assessment assumed all areas already meet the annual WHO guideline for **NO<sub>2</sub>**).

An assessment of uncertainty concludes there is a moderate degree of confidence in the modelling estimates.

---

<sup>32</sup> i.e., deaths excluding accidents and violence



## Qualitative Assessment

A known limitation of the qualitative assessment is that comparing concentrations of single contaminants with available, established ambient air quality criteria does not address additive or synergistic health effects that may occur due to exposure to mixtures of compounds.

### *Sulphur Dioxide*

Ambient levels of SO<sub>2</sub> above the WHO 10-minute guideline may have resulted in some short-term, transient effects at locations such as Whareroa Marae and Tauranga Bridge Marina including:

- Bronchoconstriction (particularly asthmatics when exercising); and/or
- Nose or throat irritation

Those most likely to experience effects would be asthmatics, babies and infants, children and elderly people. However, being transient, it is likely any effects would have been short-lived (i.e., over within minutes or hours of ambient sulphur dioxide levels returning to normal).

It is less clear what the effects of the elevated daily levels of SO<sub>2</sub> would be. A recent systemic review and meta-analysis concluded that rises in short-term SO<sub>2</sub> concentration increases the risk of all-cause mortality and respiratory mortality (Orellano *et al.*, 2021) and this was judged to have a high degree of certainty (WHO, 2021). In addition, further evidence has been reported between short-term exposure to SO<sub>2</sub> and an increased risk of asthma-associated emergency room visits and hospital admissions (Zheng *et al.*, 2021) and this relationship is considered causal (WHO, 2021).

This suggests that residents and visitors including manuhiri (guests) to Whareroa Marae and kohanga reo, residents (on boats) and visitors to the Tauranga Bridge Marina may have been, and continue to be, adversely affected by SO<sub>2</sub> emissions on occasion.

### *Hydrogen Sulphide*

Ambient air quality monitoring at Whareroa Marae and source investigation by BOPRC provides clear evidence that industrial emissions of hydrogen sulphide regularly exceeded the national guideline set to prevent against offensive odours (BOPRC, 2020). This would be reducing the quality of life and adversely impacting on the wellbeing of residents and visitors, including manuhiri.

### *Benzene*

Limited, short-term monitoring of benzene in industrial locations of the Mount Maunganui Airshed have measured ambient concentrations above a health-based, air quality criterion for acute exposure to benzene. Workers in these locations may have elevated acute exposure to benzene, however, the data are limited, and no firm conclusions can be drawn.

Whareroa Marae is within 200 metres of a bulk fuel storage facility and within 300 metres of an oil re-refining facility. Residents and visitors, including manuhiri, to Whareroa Marae and kohanga reo *may* be exposed to elevated concentrations of benzene over acute and/or chronic time frames. Benzene exposure may disproportionately impact infants and children (OEHHA, 2014). However, in the absence of any data no firm conclusions can be drawn.

There are no long-term air quality monitoring data for benzene in or around the Mount Maunganui Airshed. No conclusions can be drawn about either worker or residential chronic exposure to benzene.



### *Odour*

Odour is a well-established issue in Mount Maunganui, with more than 500 complaints to the regional council each year.

It is apparent that offensive and objectionable odours are reducing the quality of life and adversely impacting on the wellbeing of residents in and around the Mount Maunganui Airshed.

Recent literature suggests that industrial odours are associated with adverse health impacts in surrounding communities (Government of Alberta, 2017, Guadalupe-Fernandez *et al.*, 2021). This suggests that, in addition to negative impacts on wellbeing, odorous emissions may also be adversely residents' health.

### *Data Gaps*

The assessment has highlighted a lack of data for some pollutants of potential concern. Specifically, there is a dearth of ambient air quality monitoring for PAHs and benzene. This has constrained our ability to assess potential cancer risks.

## REFERENCES

---

- American Industrial Hygienists Association (AIHA), 2019. *Odor Thresholds for Chemicals with Established Occupational Health Standards*. American Industrial Hygiene Association. 2nd Edition. Falls Church. Virginia. USA.
- Agency for Toxic Substances and Disease Registry (ATSDR), 2009. *Case Studies in Environmental Medicine. Toxicity of Polycyclic Aromatic Hydrocarbons (PAHs)*. [Online: <https://www.atsdr.cdc.gov/csem/pah/docs/pah.pdf>]
- Auckland Council, 2014. *Benzene, 1,3 butadiene and other volatile organic compounds in Auckland, 2001-2013*. Technical Report 2014/037. November. Auckland. [Online: <https://knowledgeauckland.org.nz/media/1578/tr2014-037-benzene-1-3-butadiene-and-other-volatile-organic-compounds-in-auckland.pdf>]
- Bay of Plenty Regional Council, 2020. *Ambient Air Quality Data Update 2020*. Environmental Publication 2020/03. [www.boprc.govt.nz](http://www.boprc.govt.nz)
- BOPRC, undated. *Air Quality in Mount Maunganui*. [Online: <https://atlas.boprc.govt.nz/api/v1/edms/document/A3942822/content>]
- Ecocific Environmental Services, 2021. *Environmental Assessment Report. Comparison of methyl bromide and total volatile organic compounds concentrations measured at the Port of Tauranga*. Prepared for Genera Ltd. March.
- Ecocific Environmental Services, 2020. *Environmental Assessment Report. Comparison of methyl bromide and total volatile organic compounds concentrations measured at the Port of Tauranga*. Prepared for Genera Ltd. May.
- Emission Impossible Ltd, 2012. *Separation Distances for Roads*. A discussion document prepared for Auckland Council. Auckland. July.
- ESR, 2022. *Hydrogen sulphide (H<sub>2</sub>S) Factsheet*. May. [Online: <https://www.esr.cri.nz>]
- Government of Alberta, 2017. *Odours and Human Health*. Environmental Public Health Science Unit, Health Protection Branch, Public Health and Compliance Division, Alberta Health. Edmonton, Alberta. [Online: <https://open.alberta.ca/publications/9781460131534>]
- Guadalupe-Fernandez et al., 2021. Industrial Odour Pollution and Human Health: a systematic review and meta-analysis. *Environmental Health*. 20:108. doi.org/10.1186/s12940-021-00774-3.
- Hales S., Atkinson J., Metcalfe J., Kuschel G. and Woodward A., 2021. Long term exposure to air pollution mortality and morbidity in New Zealand: Cohort study. *Sci Tot Env*. Vol 801. 20 Dec. 149660. doi.org/10.1016/j.scitotenv.2021.149660
- International Agency for Research on Cancer (IARC), 2013. *IARC: Outdoor air pollution a leading environmental cause of cancer deaths*. [Online: [www.iarc.who.int](http://www.iarc.who.int)]
- Khreis H et al., 2017. Exposure to traffic-related air pollution and risk of development of childhood asthma: A systematic review and meta-analysis. *Env Int*. **100**: 1-31.

- Kuschel G., Metcalfe J., Sridhar S., Davy P., Hastings K., Mason K., Denne T., Berentson-Shaw J., Bell S., Hales S., Atkinson J., and Woodward A., 2022. *Health and Air Pollution in New Zealand 2016 (HAPINZ 3.0)*. Report prepared for Ministry for the Environment, Ministry of Health, Te Manatū Waka Ministry of Transport and Waka Kotahi NZ Transport Agency. Auckland. New Zealand. March. [Online: <https://ehinz.ac.nz>]
- Metcalfe J., and Kuschel G., *pending*. *Public Health Risks Associated with Transport Emissions in NZ: Part 2 Road Transport Emissions Trends* prepared for Ministry of Health, Ministry of Transport. Auckland. New Zealand.
- Ministry for the Environment (MfE), 2002. *Ambient Air Quality Guidelines 2002 Update*. Wellington. [Online: [www.environment.govt.nz](http://www.environment.govt.nz)]
- MfE, 2011. *2011 users' guide to the revised National Environmental Standards for Air Quality*. Updated 2014. Ministry for the Environment. Wellington. September. [Online: [www.environment.govt.nz](http://www.environment.govt.nz)]
- MfE, 2016. *Good Practice Guide for Assessing Discharges to Air from Industry*. Wellington. November. [Online: [www.environment.govt.nz](http://www.environment.govt.nz)]
- MfE, 2016b. *Good Practice Guide for Assessing and Managing Odour*. Wellington. November. [Online: [www.environment.govt.nz](http://www.environment.govt.nz)]
- MfE, 2009. *Good Practice Guide for Air Quality Monitoring and Data Management 2009*. Wellington. April. [Online: [www.environment.govt.nz](http://www.environment.govt.nz)]
- Ministry of Health (MoH), 2021a. *Mortality Collection*. Data file, Ministry of Health, Wellington, NZ. [Online: [www.health.govt.nz](http://www.health.govt.nz)]
- MoH, 2021b. *National Minimum Dataset (Hospital Events)*. Data file, Ministry of Health, Wellington, NZ. [Online: [www.health.govt.nz](http://www.health.govt.nz)]
- MoH, 2021c. *New Zealand Mortality Collection records 1996 – 2018*. 16 December. [Online]. <https://minhealthnz.shinyapps.io/mortality-web-tool/> Accessed 9 Aug 2022.
- Ministry of Transport (MoT), 2020. *Social cost of road crashes and injuries - June 2019 update*. Te Manatū Waka Ministry of Transport, Wellington, NZ
- Office of Environmental Health Hazard Assessment (OEHHA), 1999. *Air Toxics Hot Spots Program Risk Assessment Guidelines. Part 1 The Determination of Acute Reference Exposure Levels for Airborne Toxicants*. California Environmental Protection Agency. Air Toxicology and Epidemiology Section. March. [Online: <https://oehha.ca.gov/media/downloads/crn/acuterel.pdf>]
- OEHHA, 2014. *Benzene Reference Exposure Levels Technical Support Document for the Derivation of Noncancer Reference Exposure Levels. Appendix D1*. California Environmental Protection Agency. June. [Online: <https://oehha.ca.gov/media/downloads/crn/benzenerelsjune2014.pdf>]
- Orellano *et al.*, 2021. Short-term exposure to sulphur dioxide (SO<sub>2</sub>) and all-cause and respiratory mortality: A systematic review and meta-analysis. *Env Int*. Vol 150. May. 2021. 106434. <https://doi.org/10.1016/j.envint.2021.106434>

- Ostro B., 1987. Air pollution and morbidity revisited: A specification test. *J Environ Econ Manage.* **14**, 87-98.
- Pacific Air & Environment, 2009. *National Assessment of Industrial Sulphur Dioxide Emission in NZ.* Report for Ministry for the Environment. 25 March 2009.
- Public Health England (PHE), 2009. *Hydrogen Sulphide General Information.* July. [Online: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/337717/Hydrogen\\_Sulphide\\_General\\_Information\\_phe\\_v1.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/337717/Hydrogen_Sulphide_General_Information_phe_v1.pdf)]
- Sridhar S., Metcalfe J. and Mason K., 2022. *Health and Air Pollution in New Zealand 2016 (HAPINZ 3.0). He rangi hauora he iwi ora. Health effects model.* Prepared for Ministry for the Environment, Ministry of Health, Te Manatū Waka Ministry of Transport and Waka Kotahi NZ Transport Agency. Auckland. March. [Online: [www.environment.govt.nz](http://www.environment.govt.nz)]
- Office of Environmental Health Hazard Assessment (OEHHA), 2014. *Benzene Reference Exposure Levels. Technical Support Document for the Derivation of Noncancer Reference Exposure Levels. Appendix D1.* California Environmental Protection Agency. US. June. [Online: [oehha.ca.gov](http://oehha.ca.gov)]
- United States Environmental Protection Agency (US EPA), 2016. *Integrated Science Assessment for Oxides of Nitrogen—Health Criteria.* EPA/600/R-15/068. [Online: [www.federalregister.gov](http://www.federalregister.gov)]
- USEPA, 2020. *Integrated Science Assessment for Particulate Matter.* [Online: [www.epa.gov](http://www.epa.gov)]
- Waka Kotahi NZ Transport Agency, 2023. *Monetised benefits and costs manual.* April 2023. Version 1.6. Wellington. [Online: <https://www.nzta.govt.nz/assets/resources/monetised-benefits-and-costs-manual/Monetised-benefits-and-costs-manual.pdf>]
- Wickham L., Cridge B., Nicoll R., Powell J., 2021a. *Health Effects of short-term exposure to SO<sub>2</sub>.* Factsheet prepared by Emission Impossible Ltd and ESR for Ministry of Health. October. [Online: [www.esr.cri.nz](http://www.esr.cri.nz)]
- Wickham L., Cridge B., Nicoll R., Powell J., 2021b. *Effects of short-term exposure to NO<sub>2</sub>, O<sub>3</sub> and SO<sub>2</sub> on asthma.* Factsheet prepared by Emission Impossible Ltd and ESR for Ministry of Health. October. [Online: [www.esr.cri.nz](http://www.esr.cri.nz)]
- World Health Organisation (WHO), 2000. *Air Quality Guidelines for Europe.* WHO Regional Office for Europe. Copenhagen. Denmark. [Online: <https://www.who.int/>]
- WHO, 2003. *Hydrogen Sulfide: Human Health Aspects.* Geneva. Concise International Chemical Assessment Document (CICAD) NO. 53. [Online: <https://apps.who.int/iris/handle/10665/42638>]
- WHO, 2006. *Air Quality Guidelines Global Update 2005. Particulate matter, ozone, nitrogen dioxide and sulfur dioxide.* WHO Regional Office for Europe. Copenhagen. Denmark. [Online: <https://www.who.int/>]
- WHO, 2010. *Air Guidelines for Indoor Air Quality. Selected Pollutants.* WHO Regional Office for Europe. Copenhagen. Denmark. [Online: <https://www.who.int/>]

- WHO, 2013. *Review of evidence on health aspects of air pollution – REVIHAAP Project - Technical Report*. WHO Regional Office for Europe. Copenhagen. Denmark. [Online: [www.euro.who.int](http://www.euro.who.int)]
- WHO, 2014. *Health risk assessment of air pollution: General principles*. WHO Regional Office for Europe. Copenhagen. Denmark. [Online: [www.euro.who.int](http://www.euro.who.int)]
- WHO, 2016a. *Health risk assessment of air pollution. General principles*. World Health Organization Regional Office for Europe. Copenhagen. Denmark. Available at: <https://www.euro.who.int>
- WHO, 2016b. *WHO Expert Consultation: Available evidence for the future update of the WHO Global Air Quality Guidelines (AQGs)*. Meeting Report. Bonn. Germany. 29 Sep – 10 Oct 2015. WHO Regional Office for Europe. Copenhagen. Denmark. [Online: [www.euro.who.int](http://www.euro.who.int)]
- WHO, 2020. *Risk of bias assessment instrument for systematic reviews informing WHO global air quality guidelines*. Regional Office for Europe. [Online: [www.euro.who.int](http://www.euro.who.int)]
- WHO, 2021. *WHO air quality guidelines. Particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide*. Geneva: World Health Organization, 2021. [Online: <https://www.who.int/>]
- Wilton E., & Iseli J., 2019. *Tauranga Air Emissions Inventory 2018*. Prepared for Bay of Plenty Regional Council. March.
- Zheng et al., 2021. Short-term exposure to ozone, nitrogen dioxide, and sulphur dioxide and emergency department visits and hospital admissions due to asthma. A systematic review and meta-analysis. *Env Int*. Vol 150. February 2021. 106435. <https://doi.org/10.1016/j.envint.2021.106435>

# APPENDIX A: MOUNT MAUNGANUI SO<sub>2</sub> DATA

**Table A1: Summary SO<sub>2</sub> 10-minute concentrations in Mount Maunganui for 2021**

Monitoring location	Maximum 10-minute SO <sub>2</sub>	99 <sup>th</sup> percentile	Standard deviation	Exceedances WHO Guideline
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)
Rata Street	75	24	5	0
Sulphur Point	127	31	6	0
Bridge Marina	<b>1,247</b>	45	10	<b>1</b>
Whareroa Marae	361	70	13	0
Rail Yard South	82	21	4	0
Totara Street	96	29	6	0
<b>WHO 10-min Guideline SO<sub>2</sub> = 500 µg/m<sup>3</sup></b>				

**Table A2: Summary SO<sub>2</sub> 10-minute concentrations in Mount Maunganui for 2020**

Monitoring location	Maximum 10-minute SO <sub>2</sub>	99 <sup>th</sup> percentile	Standard deviation	Exceedances WHO Guideline
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)
Rata Street	80	27	6	0
Sulphur Point	142	35	7	0
Bridge Marina	275	44	10	0
Whareroa Marae	432	72	15	0
Rail Yard South	109	22	5	0
Totara Street	90	34	6	0
<b>WHO 10-min Guideline SO<sub>2</sub> = 500 µg/m<sup>3</sup></b>				

**Table A3: Summary SO<sub>2</sub> 10-minute concentrations in Mount Maunganui for 2019**

Monitoring location	Maximum 10-minute SO <sub>2</sub>	99 <sup>th</sup> percentile	Standard deviation	Exceedances WHO Guideline
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)
Rata Street	<b>775</b>	197	42	<b>10</b>
Sulphur Point	287	106	19	0
Bridge Marina	232	76	15	0
Whareroa Marae	472	113	23	0
Rail Yard South	393	154	32	0
Totara Street	359	85	19	0
<b>WHO 10-min Guideline SO<sub>2</sub> = 500 µg/m<sup>3</sup></b>				

**Table A4: Summary SO<sub>2</sub> 1-hour concentrations in Mount Maunganui for 2021**

Monitoring location	Maximum 1-hour SO <sub>2</sub>	99 <sup>th</sup> percentile	Exceedances NES lower limit (350 µg/m <sup>3</sup> )	Exceedances NES upper limit (570 µg/m <sup>3</sup> )
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)	(no.)
Rata Street	63	23	0	0
Sulphur Point	96	29	0	0
Bridge Marina	312	40	0	0
Whareroa Marae	125	59	0	0
Rail Yard South	38	18	0	0
Totara Street	60	27	0	0

**Table A5: Summary SO<sub>2</sub> 1-hour concentrations in Mount Maunganui for 2020**

Monitoring location	Maximum 1-hour SO <sub>2</sub>	99 <sup>th</sup> percentile	Exceedances NES lower limit (350 µg/m <sup>3</sup> )	Exceedances NES upper limit (570 µg/m <sup>3</sup> )
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)	(no.)
Rata Street	57	25	0	0
Sulphur Point	100	33	0	0
Bridge Marina	161	40	0	0
Whareroa Marae	251	60	0	0
Rail Yard South	68	20	0	0
Totara Street	64	29	0	0

**Table A6: Summary SO<sub>2</sub> 1-hour concentrations in Mount Maunganui for 2019**

Monitoring location	Maximum 1-hour SO <sub>2</sub>	99 <sup>th</sup> percentile	Exceedances NES lower limit (350 µg/m <sup>3</sup> )	Exceedances NES upper limit (570 µg/m <sup>3</sup> )
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)	(no.)
Rata Street	<b>575</b>	173	<b>4</b>	<b>1</b>
Sulphur Point	208	96	0	0
Bridge Marina	157	65	0	0
Whareroa Marae	206	97	0	0
Rail Yard South	226	126	0	0
Totara Street	167	72	0	0



**Table A7: Summary SO<sub>2</sub> daily concentrations in Mount Maunganui for 2021**

Monitoring location	Maximum daily SO <sub>2</sub>	99% percentile	Standard deviation	Exceedances WHO 2021 AQG
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)
Rata Street	15	13	3	0
Sulphur Point	39	19	4	0
Bridge Marina	36	24	5	0
Whareroa Marae	<b>42</b>	24	6	<b>1</b>
Rail Yard South	12	11	2	0
Totara Street	26	14	3	0
<b>WHO 2021 AQG Daily SO<sub>2</sub> = 40 µg/m<sup>3</sup></b>				

**Table A8: Summary SO<sub>2</sub> daily concentrations in Mount Maunganui for 2020**

Monitoring location	Maximum daily SO <sub>2</sub>	99% percentile	Standard deviation	Exceedances WHO 2021 AQG
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)
Rata Street	16	13	3	0
Sulphur Point	24	18	4	0
Bridge Marina	<b>54</b>	29	6	<b>3</b>
Whareroa Marae	<b>54</b>	28	6	<b>1</b>
Rail Yard South	17	12	2	0
Totara Street	19	15	3	0
<b>WHO 2021 AQG Daily SO<sub>2</sub> = 40 µg/m<sup>3</sup></b>				

**Table A9: Summary SO<sub>2</sub> daily concentrations in Mount Maunganui for 2019**

Monitoring location	Maximum daily SO <sub>2</sub>	99% percentile	Standard deviation	Exceedances WHO 2021 AQG
	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(µg/m <sup>3</sup> )	(no.)
Rata Street	<b>140</b>	<b>90</b>	21	<b>50</b>
Sulphur Point	<b>93</b>	<b>60</b>	12	<b>11</b>
Bridge Marina	<b>44</b>	33	7	<b>1</b>
Whareroa Marae	<b>48</b>	<b>42</b>	10	<b>5</b>
Rail Yard South	<b>92</b>	<b>72</b>	18	<b>62</b>
Totara Street	<b>54</b>	<b>40</b>	9	<b>4</b>
<b>WHO 2021 AQG Daily SO<sub>2</sub> = 40 µg/m<sup>3</sup></b>				




Figure A1: 10-minute SO<sub>2</sub> at Rata Street and Sulphur Point: 2019 – 2021







Figure A2: 10-minute SO<sub>2</sub> at Rail Yard South and Totara Street: 2019 – 2021




**Figure A3: 10-minute SO<sub>2</sub> at Whareroa Marae and Tauranga Bridge Marina: 2019 – 2021**



**Figure A4: Hourly SO<sub>2</sub> at Rata Street and Sulphur Point: 2019 – 2021**



**Figure A5: Hourly SO<sub>2</sub> at Rail Yard South and Totara Street: 2019 – 2021**



**Figure A6: Hourly SO<sub>2</sub> at Whareroa Marae and Tauranga Bridge Marina: 2019 – 2021**




Figure A7: Daily SO<sub>2</sub> at Rata Street and Sulphur Point: 2019 – 2021




Figure A8: Daily SO<sub>2</sub> at Rail Yard South and Totara Street: 2019 and 2021




Figure A9: Daily SO<sub>2</sub> at Whareroa Marae and Tauranga Bridge Marina: 2019 – 2021




Figure A10: Frequency of (10-minute) wind direction (from) and wind speed measured at Whareroa Marae; 1 Jan – 31 Dec 2019






Figure A11: Frequency of (10-minute) wind direction (from) and wind speed measured at Whareroa Marae; 1 Jan – 31 Dec 2020




Figure A12: Frequency of (10-minute) wind direction (from) and wind speed measured at Whareroa Marae; 1 Jan – 31 Dec 2021



**INSTITUTE OF ENVIRONMENTAL  
SCIENCE AND RESEARCH LIMITED**

▀ **Kenepuru Science Centre**  
34 Kenepuru Drive, Kenepuru, Porirua 5022  
PO Box 50348, Porirua 5240  
New Zealand  
T: +64 4 914 0700 F: +64 4 914 0770

▀ **Mt Albert Science Centre**  
120 Mt Albert Road, Sandringham, Auckland 1025  
Private Bag 92021, Auckland 1142  
New Zealand  
T: +64 9 815 3670 F: +64 9 849 6046

▀ **NCBID – Wallaceville**  
66 Ward Street, Wallaceville, Upper Hutt 5018  
PO Box 40158, Upper Hutt 5140  
New Zealand  
T: +64 4 529 0600 F: +64 4 529 0601

▀ **Christchurch Science Centre**  
27 Creyke Road, Ilam, Christchurch 8041  
PO Box 29181, Christchurch 8540  
New Zealand  
T: +64 3 351 6019 F: +64 3 351 0010

[www.esr.cri.nz](http://www.esr.cri.nz)